

TopBP1 (D8G4L) Rabbit mAb

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W, IP	H	Endogenous	170	Rabbit IgG	#Q92547	11073

Product Usage Information**Application**

Western Blotting
Immunoprecipitation

Dilution

1:1000
1:100

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. *Do not aliquot the antibody.*

Specificity/Sensitivity

TopBP1 (D8G4L) Rabbit mAb recognizes endogenous levels of total TopBP1 protein.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Leu27 of human TopBP1 protein.

Background

Topoisomerase II Binding Protein 1 (TopBP1) contains eight BRCT domains, which facilitate interaction with various proteins, phosphopeptides, and DNA. Through these interactions, TopBP1 functions in the regulation of DNA replication, DNA repair, checkpoint control, and transcription (1). TopBP1 contacts the checkpoint kinase ATR and its binding partner ATRIP, and induces ATR and Chk1 activation in collaboration with claspin (2,3). Activation of ATR is dependent on recruitment of TopBP1 through the MRN (MRE11-RAD50-NBS1) complex, and the 911 (RAD9-RAD1-HUS1) complex is required for full activation (4). TopBP1 stabilizes Bloom syndrome helicase (BLM) during S phase of the cell cycle to suppress sister chromatin exchange and maintain genome stability (5). TopBP1 also regulates initiation of DNA replication along with the DNA replication factor treslin (6,7). TopBP1 has been shown to prevent replication associated DNA damage during neurogenesis (8), and to interact with mutant p53, mediating mutant p53 gain-of-function activity such as growth promotion and resistance to chemotherapeutic drugs (9). Phosphorylation of TopBP1 at Ser1159 by Akt regulates TopBP1 oligomerization and function in E2F1-dependent transcriptional regulation (10).

Background References

- Garcia, V. et al. (2005) *DNA Repair (Amst)* 4, 1227-39.
- Lindsey-Boltz, L.A. and Sancar, A. (2011) *J Biol Chem* 286, 19229-36.
- Burrows, A.E. and Elledge, S.J. (2008) *Genes Dev* 22, 1416-21.
- Duursma, A.M. et al. (2013) *Mol Cell* 50, 116-22.
- Wang, J. et al. (2013) *Mol Cell* 52, 667-78.
- Kumagai, A. et al. (2010) *Cell* 140, 349-59.
- Boos, D. et al. (2013) *Science* 340, 981-4.
- Lee, Y. et al. (2012) *Nat Neurosci* 15, 819-26.
- Liu, K. et al. (2011) *Mol Cell Biol* 31, 4464-81.
- Liu, K. et al. (2013) *Mol Cell Biol* 33, 4685-700.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IP:** Immunoprecipitation

Cross-Reactivity Key

H: Human

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.