

## Phospho-MAPK Substrates Motif [PXpTP] MultiMab<sup>®</sup> Rabbit mAb mix



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W	Reactivity:	Sensitivity:	Source/Isotype:
VV	All	Endogenous	Rabbit IgG
Product Usage Information		<b>Application</b> Western Blotting	<b>Dilution</b> 1:1000
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 $\mu$ g/ml BSA and 50% glycerol. Store at – 20°C. Do not aliquot the antibody.	
Specificity/Sensitivity		Phospho-MAPK Substrates Motif [PXpTP] MultiMab <sup>®</sup> Rabbit mAb mix recognizes endogenous levels of proteins that are phosphorylated at threonine within the context of a PXTP motif. The antibody does not cross-react with endogenous levels of non-phosphorylated proteins, phospho-threonine in another context, or proteins with a phosphorylated serine within the PXSP motif.	
Source / Purification		MultiMab <sup>®</sup> rabbit monoclonal mix antibodies are prepared by combining individual rabbit monoclonal clones in optimized ratios for the approved applications. Each antibody in the mix is carefully selected based on motif recognition and performance in multiple assays. Each mix is engineered to yield the broadest possible coverage of the modification being studied while ensuring a high degree of specificity for the modification or motif.	
Background		The MAPK and CDK families of serine/threonine protein kinases play important roles in proliferation and cell cycle control. These kinases phosphorylate threonine or serine followed by a proline residue (1-3). MAPK phosphorylates substrates with the consensus sequence PX(S/T)P, and CDKs phosphorylate substrates containing the consensus sequence (S/T)PXR/K. Cell Signaling Technology has developed antibodies that bind to phospho-threonine followed by proline, motifs PXS*/T*P and/or S*PXR/K, for use in the study and discovery of new MAPK and CDK substrates (4,5).	
Background References		1. Cross, T.G. et al. (2000) <i>Exp Cell Res</i> 256, 34-41. 2. Reynolds, C.H. et al. (2000) <i>J Neurochem</i> 74, 1587-95. 3. Seger, R. and Krebs, E.G. (1995) <i>FASEB J</i> 9, 726-35. 4. Holmes, J.K. and Solomon, M.J. (1996) <i>J Biol Chem</i> 271, 25240-6. 5. Songyang, Z. et al. (1996) <i>Mol Cell Biol</i> 16, 6486-93.	

**Species Reactivity** Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X

TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key W: Western Blotting

Cross-Reactivity Key All: All Species Expected

**Trademarks and Patents** Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

MultiMab is a registered trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for

more information.

**Limited Uses** Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST,

the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no

force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.