


Revision 3

Store at
-20C
#14434

SF3B1 (D7L5T) Rabbit mAb



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W, W-S	H M R Mk	Endogenous	155	Rabbit IgG	#O75533	23451

Product Usage Information

Application

Western Blotting
Simple Western™

Dilution

1:1000
1:10 - 1:50

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. *Do not aliquot the antibody.*

Specificity/Sensitivity

SF3B1 (D7L5T) Rabbit mAb recognizes endogenous levels of total SF3B1 protein.

Species predicted to react based on 100% sequence homology

Hamster, Bovine

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues near the amino terminus of human SF3B1 protein.

Background

Splicing factor 3b subunit 1 (SF3B1) is an integral component of the U2 small nuclear ribonucleoprotein (U2 snRNP) and plays an important role in the splicing of pre-mRNA that involves the removal of introns and the joining of exons to form mature mRNA (1-3). The assembly and proper recognition of splice sites are driven by sequences at the pre-mRNA intron-exon splice sites. The 5' splice donor site is recognized by the U1 snRNP complex, while U2 snRNP binds to the 3' splice site (branch point), ensuring the anchoring of the spliceosome machinery at the splice sites (3,4). Recent whole exome sequencing studies have demonstrated a high incidence of somatic mutations of *SF3B1* in patients with various hematological malignancies such as chronic lymphocytic leukemia and myelodysplastic syndromes (2,3,5,6). Misregulation of pre-mRNA splicing arising from mutations of the spliceosome components such as SF3B1 is thought to contribute to changes in the expression patterns of key proteins that are involved in pathways such as cell cycle progression, cell death, and cancer metabolism (2,3).

Background References

1. Jurica, M.S. and Moore, M.J. (2003) *Mol Cell* 12, 5-14.
2. Cazzola, M. et al. (2013) *Blood* 121, 260-9.
3. Bonnal, S. et al. (2012) *Nat Rev Drug Discov* 11, 847-59.
4. Gozani, O. et al. (1998) *Mol Cell Biol* 18, 4752-60.
5. Quesada, V. et al. (2012) *Nat Genet* 44, 47-52.
6. Baliakas, P. et al. (2015) *Leukemia* 29, 329-36.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **W-S:** Simple Western™

Cross-Reactivity Key

H: Human **M:** Mouse **R:** Rat **Mk:** Monkey

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.

Update your browser for more security, speed and the best experience on this site.

Update browser

Orders: 877-616-CELL (2355) • orders@cellsignal.com • Support: 877-678-TECH (8324) • info@cellsignal.com • Web: cellsignal.com
For Research Use Only. Not for Use in Diagnostic Procedures.