

CLASP2 (D5K3E) Rabbit mAb

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W	H M R Mk	Endogenous	165	Rabbit IgG	#O75122	23122

Product Usage Information**Application**

Western Blotting

Dilution

1:1000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. *Do not aliquot the antibody.*

Specificity/Sensitivity

CLASP2 (D5K3E) Rabbit mAb recognizes endogenous levels of total CLASP2 protein. This antibody also cross-reacts with an unidentified protein of 95 kDa.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Ala289 of human CLASP2 protein.

Background

Microtubules (MTs) are polarized cellular filaments composed of α/β tubulin heterodimers. The slower growing (minus) microtubule ends are located at MT organizing centers (MTOCs), with the faster growing (plus) ends extending to the cell periphery. The regulation of MT dynamics is an important part of several biological processes, including cell division, migration, adhesion, membrane trafficking, and polarity (1).

Human cytoplasmic linker-associate proteins 1 and 2 (CLASP1 and CLASP2) are evolutionarily conserved proteins that localize to the plus ends of interphase microtubules. During mitosis, CLASP 1 and CLASP2 localize to the centrosomes and kinetochores (KT) where they regulate mitotic spindle positioning to ensure proper chromosome alignment (2,3). Research studies indicate that phosphorylation of the carboxy terminus of CLASP2 during mitosis by CDK1 and PLK1 is required for efficient mitotic MT-KT attachment (4). Phosphorylation of CLASP2 at Ser1013 is a critical step that primes CLASP2 for further phosphorylation by PLK1 (4). The additional phosphorylation of CLASP2 at Ser533 and Ser537 by GSK3- β controls the distribution of CLASP2 on MTs by inhibiting CLASP2 interaction with the Rac1/cdc42 effector protein IQGAP1 (5).

Background References

1. Wiese, C. and Zheng, Y. (2006) *J Cell Sci* 119, 4143-53.
2. Mimori-Kiyosue, Y. et al. (2005) *J Cell Biol* 168, 141-53.
3. Logarinho, E. et al. (2012) *Nat Cell Biol* 14, 295-303.
4. Maia, A.R. et al. (2012) *J Cell Biol* 199, 285-301.
5. Watanabe, T. et al. (2009) *J Cell Sci* 122, 2969-79.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting

Cross-Reactivity Key

H: Human **M:** Mouse **R:** Rat **Mk:** Monkey

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.