

Store at
-20°C

#14708

Anti-rabbit IgG (H+L), Biotinylated Antibody

www.cellsignal.com

Support: 877-678-TECH (8324)
www.cellsignal.com/support

Orders: 877-616-CELL (2355)
orders@cellsignal.com

rev. 02/07/17

For Research Use Only. Not For Use In Diagnostic Procedures.

Description: Affinity purified goat anti-rabbit IgG (H+L) antibody is conjugated to biotin. This product has been optimized for use as a secondary antibody in western blotting applications.

Background: Biotinylated anti-species IgG conjugated antibodies enable signal amplification and are ideal for detecting lower expressing proteins.

Specificity/Sensitivity: Anti-rabbit IgG (H+L), Biotinylated Antibody reacts with the heavy and light chains of most rabbit immunoglobulins. Less than 1% cross-reactivity is observed with immunoglobulins from other species.

Storage: Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. *Do not aliquot the antibody.*

Recommended Antibody Dilutions:

Western blotting 1:1000 – 1:3000

For product specific protocols and a complete listing of recommended companion products please see the product web page at www.cellsignal.com

Thank you for your recent purchase. If you would like to provide a review visit www.cellsignal.com/comments.

© 2014 Cell Signaling Technology, Inc.
Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.



Cell Signaling
TECHNOLOGY®

Applications: W—Western IP—Immunoprecipitation IHC—Immunohistochemistry ChIP—Chromatin Immunoprecipitation IF—Immunofluorescence F—Flow cytometry E-P—ELISA-Peptide **Species Cross-Reactivity:** H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken Dm—D. melanogaster X—Xenopus Z—zebrafish B—bovine Dg—dog Pg—pig Sc—S. cerevisiae Ce—C. elegans Hr—Horse All—all species expected Species enclosed in parentheses are predicted to react based on 100% homology.