

<b>#14774</b>	<b>Store at RT</b>	<h1>Chloroquine</h1>	 <b>Orders:</b> 877-616-CELL (2355) orders@cellsignal.com <b>Support:</b> 877-678-TECH (8324) <b>Web:</b> info@cellsignal.com cellsignal.com
	150 mg	<b>3 Trask Lane   Danvers   Massachusetts   01923   USA</b>	

**For Research Use Only. Not for Use in Diagnostic Procedures.**

### Background

Chloroquine (CQ) is a lysosomotropic agent with an extensive range of biological effects (1). Historically known for its anti-malarial activity, chloroquine is a widely used biological research tool for studying autophagy inhibition. Research studies demonstrate that chloroquine accumulates in acidic lysosomes and increases the lysosomal pH. This inhibits lysosomal hydrolases and prevents autophagosomal fusion and degradation, which can result in apoptotic or necrotic cell death (1-4). Inhibition of chloroquine-induced apoptosis with the V-ATPase inhibitor bafilomycin A1 has been observed in several cell types (4). Chloroquine also enhances the anti-neoplastic effects of the histone deacetylase inhibitor vorinostat (SAHA) (5).

Chloroquine treatment of cells leads to accumulation of light chain 3-II (LC3-II) (1-3). This autophagy marker resides within autophagosomal membranes during the autophagic process and is degraded upon fusion with lysosomes. Chloroquine inhibition of these fusion events effectively blocks LC3-II degradation.

This product has applications to SARS-CoV-2 research into the mechanisms of the Novel Coronavirus, which has caused the COVID-19 pandemic.

### Molecular Formula

C<sub>18</sub>H<sub>26</sub>ClN<sub>3</sub>•2H<sub>3</sub>PO<sub>4</sub>

### Molecular Weight

515.9 g/mol

### Purity

>98%

### CAS

50-63-5

### Solubility

Soluble in H<sub>2</sub>O at 25mg/ml.

### Storage

Store lyophilized at room temperature or in solution at -20°C, desiccated. Protect from light. In lyophilized form, the chemical is stable for 24 months. Once in solution, use within 3 months to prevent loss of potency. Aliquot to avoid multiple freeze/thaw cycles.

### Directions for Use:

Chloroquine is supplied as a lyophilized powder. For a 50 mM stock, reconstitute the 150 mg in 5.82 ml sterile dH<sub>2</sub>O. First add 1 ml dH<sub>2</sub>O to the tube containing the chemical, vortex, and dispense into a new, larger tube. Repeat this action two or three more times to transfer any residual material. Add additional dH<sub>2</sub>O to the new tube to bring the volume up to 5.82 ml. Filter sterilize into sterile tube. Utilize a syringe and 0.2 µm syringe filter to minimize sample loss.

Working concentrations and length of treatment can vary depending on the desired effect, but it is typically used at 25-100 µM for 12-48 hr.

### Background References

1. Liang, X. et al. (2012) *Cancer Res* 72, 2791-801.
2. Liang, X. et al. (2014) *Cell Biosci* 4, 10.
3. Kung, C.P. et al. (2011) *Crit Rev Eukaryot Gene Expr* 21, 71-100.
4. Shacka, J.J. et al. (2006) *Mol Pharmacol* 69, 1125-36.
5. Carew, J.S. et al. (2007) *Blood* 110, 313-22.

### Cross-Reactivity Key

**H:** human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster  
**X:** Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse  
**GP:** Guinea Pig **Rab:** rabbit **All:** all species expected

### Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.  
Alexa Fluor is a registered trademark of Life Technologies Corporation.  
All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

**Limited Uses**

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.