

TFE3 Antibody



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W	Reactivity: H M R	Sensitivity: Endogenous	MW (kDa): 72-89	Source/Isotype: Rabbit	UniProt ID: #P19532	Entrez-Gene Id: 7030
---------------------------	-----------------------------	-----------------------------------	---------------------------	----------------------------------	-------------------------------	--------------------------------

Product Usage Information

Application

Western Blotting

Dilution

1:1000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

TFE3 Antibody recognizes endogenous levels of total TFE3 protein. The upper band is presumed to be sumoylated TFE3 protein. This antibody does not cross react with the closely related MITF protein.

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues near the carboxy terminus of human TFE3 protein. Antibodies are purified by protein A and peptide affinity chromatography.

Background

Transcription factor E3 (TFE3) is a member of a family of basic helix-loop-helix leucine zipper transcription factors that include MITF, TFEB, TFE3, and TFEC. Members of this family form heterodimers with each other, bind the same DNA sequences, and undergo the same types of post-translational modifications, including sumoylation (1). Research studies indicate that TFE3 and other family members play roles in development, organelle biogenesis, nutrient sensing, autophagy, and energy metabolism (2,3). Additional studies report that TFE3 controls the gate for pluripotent cells to exit the state of pluripotency prior to differentiation (4). Translocations involving the *TFE3* gene region have been identified in a number of tumors, including sporadic renal cell tumors. Several specific translocations that result in kidney cancer and involve the *TFE3* gene have been described and characterized in detail (5).

Background References

1. Miller, A.J. et al. (2005) *J Biol Chem* 280, 146-55.
2. Martina, J.A. et al. (2014) *Cell Mol Life Sci* 71, 2483-97.
3. Martina, J.A. et al. (2014) *Sci Signal* 7, ra9.
4. Betschinger, J. et al. (2013) *Cell* 153, 335-47.
5. Kauffman, E.C. et al. (2014) *Nat Rev Urol* 11, 465-75.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting

Cross-Reactivity Key

H: Human **M:** Mouse **R:** Rat

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in

any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.