#

-

## NF-кB2 p100/p52 (18D10) Rabbit mAb (PE Conjugate) 2005



Orders:	877-616-CELL (2355) orders@cellsignal.com
Support:	877-678-TECH (8324)
Web:	info@cellsignal.com cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: FC-FP	<b>Reactivity:</b> H Mk	<b>Sensitivity:</b> Endogenous	<b>Source/Isotype:</b> Rabbit IgG	<b>UniProt ID:</b> #Q00653	Entrez-Gene Id: 4791		
Product Usage Information		<b>Application</b> Flow Cytometry (Fixed/Permeabilized)		<b>Dilution</b> 1:50			
Storage		Supplied in PBS (pH 7.2), less than 0.1% sodium azide and 2 mg/ml BSA. Store at 4°C. Do not aliquot the antibodies. Protect from light. Do not freeze.					
Specificity/Sensi	tivity	NF-κB2 p100/p52 (18D10) Rabbit mAb (PE Conjugate) recognizes endogenous levels of both the p100 precursor and the p52 protein active form of NF-κB2. The antibody does not cross-react with other family members.					
Source / Purifica	tion	Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues at the amino-terminus of human NF-кB2 p100/p52 protein.					
Description		This Cell Signaling Technology antibody is conjugated to phycoerythrin (PE) and tested in-house for direct flow cytometry analysis in human cells. The antibody is expected to exhibit the same species cross-reactivity as the unconjugated NF-κB2 p100/p52 (18D10) Rabbit mAb #3017.					
Background		Transcription factors of the nuclear factor κB (NF-κB)/Rel family play a pivotal role in inflammatory and immune responses (1,2). There are five family members in mammals: RelA, c-Rel, RelB, NF-κB1 (p105/p50), and NF-κB2 (p100/p52). Both p105 and p100 are proteolytically processed by the proteasome to produce p50 and p52, respectively. Rel proteins bind p50 and p52 to form dimeric complexes that bind DNA and regulate transcription. In unstimulated cells, NF-κB is sequestered in the cytoplasm by IkB inhibitory proteins (3-5). NF-κB-activating agents can induce the phosphorylation of IkB proteins, targeting them for rapid degradation through the ubiquitin-proteasome pathway and releasing NF-κB to enter the nucleus where it regulates gene expression (6-8). NIK and IKKα (IKK1) regulate the phosphorylation and processing of NF-κB2 (p100) to produce p52, which translocates to the nucleus (9-11).					
Background Ref	erences	<ol> <li>Baeuerle, P.A. and Henkel, T. (1994) Annu Rev Immunol 12, 141-79.</li> <li>Baeuerle, P.A. and Baltimore, D. (1996) Cell 87, 13-20.</li> <li>Haskill, S. et al. (1991) Cell 65, 1281-9.</li> <li>Thompson, J.E. et al. (1995) Cell 80, 573-82.</li> <li>Whiteside, S.T. et al. (1997) EMBO J 16, 1413-26.</li> <li>Traenckner, E.B. et al. (1995) EMBO J 14, 2876-83.</li> <li>Scherer, D.C. et al. (1995) Proc Natl Acad Sci USA 92, 11259-63.</li> <li>Chen, Z.J. et al. (1996) Cell 84, 853-62.</li> <li>Senftleben, U. et al. (2001) Science 293, 1495-9.</li> <li>Coope, H.J. et al. (2002) EMBO J 21, 5375-85.</li> <li>Xiao, G. et al. (2001) Mol Cell 7, 401-9.</li> </ol>					
Species Reactivi	ty	Species reactivity is determined by testing in at least one approved application (e.g., western blot).					
Applications Key	,	FC-FP: Flow Cytometry (Fixed/Permeabilized)					
Cross-Reactivity	Кеу	H: Human Mk: Monkey					
Trademarks and	Patents	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.					
		XP is a registered trademark of Cell Signaling Technology, Inc.					
		All other trademarks are more information.	the property of their res	pective owners. Visit	cellsignal.com/trademarks for		

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.