His-Tag (D3I1O) XP[®] Rabbit mAb (PE Conjugate)



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: FC-FP	Reactivity: All	Sensitivity: Source/Isotype: Transfected Only Rabbit IgG		
Product Usage Information		Application Flow Cytometry (Fixed/Permeabilized)	Dilution 1:50	
Storage		Supplied in PBS (pH 7.2), less than 0.1% sodium azide and 2 mg/ml BSA. Store at 4°C. Do not aliquot the antibodies. Protect from light. Do not freeze.		
Specificity/Sensitivity		His-Tag (D3I1O) XP [®] Rabbit mAb (PE Conjugate) recognizes recombinant proteins containing the 6xHis epitope tag. The antibody recognizes the 6xHis-tag fused to either the amino or carboxy terminus of targeted proteins in transfected cells.		
Source / Purification		Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues of the 6xHis epitope tag.		
Description		This Cell Signaling Technology antibody is conjugated to phycoerythrin (PE) and tested in-house for direct flow cytometry analysis in human cells. The antibody is expected to exhibit the same species cross-reactivity as the unconjugated His-Tag (D3I1O) XP [®] Rabbit mAb #12698.		
Background		Epitope tags are useful for the labeling and detection of proteins using immunoblotting, immunoprecipitation, and immunostaining techniques. Because of their small size, they are unlikely to affect the tagged protein's biochemical properties.		
		A variety of plasmids contain DNA that encodes an amino-terminal ta residues followed by an extended multiple cloning site. The 6xHis tag proteins allows for efficient coupling to Ni ²⁺ affinity resins and purific chromatography (1).	on the expressed recombinant	
		As is the case with other protein tag systems (2), this polyhistidine tag can often be cleaved at sites recognized by proteases such as thrombin and enterokinases to isolate the protein of interest (1).		
Background References		1. Kroll, D.J. et al. (1993) <i>DNA Cell Biol</i> 12, 441-53. 2. di Guan, C. et al. (1988) <i>Gene</i> 67, 21-30.		
Species Reactivity		Species reactivity is determined by testing in at least one approved ap	oplication (e.g., western blot).	
Applications Key	ations Key FC-FP: Flow Cytometry (Fixed/Permeabilized)			
Cross-Reactivity K	ey	All: All Species Expected		
Trademarks and Patents		Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.		
		XP is a registered trademark of Cell Signaling Technology, Inc.		
		All other trademarks are the property of their respective owners. Visit more information.	cellsignal.com/trademarks for	
Limited Uses		Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.		
		Products are labeled with For Research Use Only or a similar labeling		

approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in

any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.