

# Phospho-Autocam Biotinylated Peptide

✓ 1.25 ml at 6 µM



Cell Signaling  
TECHNOLOGY®

**Orders** ■ 877-616-CELL (2355)  
orders@cellsignal.com

**Support** ■ 877-678-TECH (8324)  
info@cellsignal.com

**Web** ■ www.cellsignal.com

New 04/08

This product is for *in vitro* research use only and is not intended for use in humans or animals.

**Description:** Autocam is a synthetic peptide derived from the Ca<sup>2+</sup>/Calmodulin-Dependent Protein Kinase II (CaMKII) autophosphorylation site that serves as a selective CaMKII substrate. The threonine residue in the peptide has been chemically phosphorylated in the course of peptide synthesis. This phosphopeptide was generated for use as a positive control in kinase assays (CaMKI-γ Kinase #7456, MAPKAPK-2 Kinase #7442) but it may also serve as a positive control in other heterogeneous or homogeneous kinase assays.

**Peptide Core Sequence:** RQET\*VDA

**Molecular Weight:** 2088 daltons

**Quality Control:** The quality of the biotinylated peptide was evaluated by reverse-phase HPLC and by mass spectrometry.

**Directions for Use:** This phosphorylated peptide can be detected with the Phospho-Threonine (42H4) Mouse mAb #9386. A sample kinase assay protocol is attached.

**Storage:** Supplied in 0.0001% DMSO. Store at -20°C.

**Companion Products:**

CaMKI-γ Kinase #7456

MAPKAPK-2 Kinase #7442

Phospho-Threonine (42H4) Mouse mAb #9386

Autocam Biotinylated Peptide #1411

## Protocol for Serine/Threonine Kinase Assay

**IMPORTANT:** Use of an automated microplate washer as well as centrifugation of plates when appropriate, greatly improves reproducibility.

### A Additional Solutions and Reagents (Not included)

1. **Wash Buffer:** 1X PBS, 0.05% Tween-20 (PBS/T)
2. Bovine Serum Albumin (BSA)
3. **Stop Buffer:** 50 mM EDTA pH 8
4. Kinase Buffer (10X) #9802
5. ATP (10 mM) #9804
6. Active kinase (See companion products)
7. Primary antibody (See companion products)

### B Suggested Protocol for 100 Assays

1. Add 100 µl 10 mM ATP to 1.25 ml 6-12 µM substrate peptide. Adjust the mixture with dH<sub>2</sub>O to 2.5 ml to make 2X ATP/substrate cocktail ([ATP]=400 µM, [substrate] = 3-6 µM).
2. Transfer enzyme from -80°C to ice. Allow enzyme to thaw on ice.
3. **Microcentrifuge briefly at 4°C to bring liquid to the bottom of the vial. Return immediately to ice.**
4. Add 1 ml 10X kinase buffer [250 mM Tris-HCl pH 7.5, 100 mM MgCl<sub>2</sub>, 1 mM Na<sub>3</sub>VO<sub>4</sub>, 50 mM β-glycerophosphate, 20 mM dithiothreitol (DTT)] to 1.5 ml dH<sub>2</sub>O to make 2.5 ml 4X reaction buffer.
5. Dilute enzyme in 1.25 ml of 4X reaction buffer to make 4X reaction cocktail ([enzyme]=0.8-8.0 ng/µl in 4X reaction cocktail).
6. Add 12.5 µl of the 4X reaction cocktail to 12.5 µl/well of prediluted compound of interest (usually around 10 µM) and incubate for 5 minutes at room temperature.
7. Add 25 µl of 2X ATP/substrate cocktail to 25 µl/well preincubated reaction cocktail/compound.

#### Final Assay Conditions for a 50 µl Reaction

25 mM Tris-HCl (pH 7.5)  
10 mM MgCl<sub>2</sub>  
5 mM β-glycerophosphate  
0.1 mM Na<sub>3</sub>VO<sub>4</sub>  
2 mM DTT  
200 µM ATP  
1.5-3 µM peptide  
10-100 ng kinase

8. Incubate reaction plate at room temperature for 30 minutes.
9. Add 50 µl/well Stop Buffer (50 mM EDTA, pH 8) to stop the reaction.
10. Transfer 25 µl of each reaction to a 96-well streptavidin-coated plate containing 75 µl dH<sub>2</sub>O/well and incubate at room temperature for 60 minutes.
11. Wash three times with 200 µl/well PBS/T.
12. Dilute primary antibody in PBS/T with 1% BSA. \*Add 100 µl/well primary antibody (1:500 dilution for mouse mAb or 1:1000 dilution for rabbit mAb or polyclonal antibody)
13. Incubate at 37°C for 120 minutes.
14. Wash three times with 200 µl/well PBS/T.
15. For DELFIA® or Colorimetric ELISA detection methods please use the following protocols.

DELFIA® is a registered trademark of PerkinElmer Life Sciences

### DELFIA® Assay

1. Prepare appropriate dilution of Europium labeled secondary antibody in PBS/T with 1% BSA (1:500 dilution for anti-mouse IgG or 1:1000 for anti-rabbit IgG).
  2. Add 100 µl/well secondary antibody solution.
  3. Incubate at room temperature for 30 minutes.
  4. \*Wash five times with 200 µl/well PBS/T.
  5. Add 100 µl/well DELFIA® Enhancement Solution.
  6. Incubate at room temperature for 5 minutes.
  7. Read plate using a Time Resolved Fluorescent plate reader using the following settings;
    - a. Excitation Filter: 340 nm
    - b. Emission Filter: 615 nm
    - c. Delay\*\*: 400 µs
- \*\* Delay time is the delay from the excitation pulse to the beginning of the measurement.

### Companion Products for DELFIA®

DELFIA® Europium-labeled Anti-mouse IgG (PerkinElmer Life Sciences #AD0124)  
DELFIA® Europium-labeled Anti-rabbit IgG (PerkinElmer Life Sciences #AD0105)  
DELFIA® Enhancement Solution (PerkinElmer Life Sciences #1244-105)  
DELFIA® Streptavidin coated, 96-well, yellow plate (PerkinElmer Life Sciences AAAND-0005)

### Colorimetric ELISA Assay

1. Prepare appropriate dilution of HRP labeled secondary antibody in PBS/T with 1% BSA (1:500 dilution for anti-mouse IgG or 1:1000 for anti-rabbit IgG).
2. Add 100 µl/well secondary antibody solution.
3. Incubate at room temperature for 30 minutes.
4. \*Wash five times with 200 µl/well PBS/T.
5. Add 100 µl/well TMB substrate.
6. Incubate at room temperature for 15 minutes.
7. Add 100 µl/well of stop solution.
8. Mix well.
9. Read the absorbance at 450 nm with a microtiter plate reader.

### Companion Products For Colorimetric ELISA Assay

Anti-mouse IgG, HRP Linked Antibody #7076  
Anti-rabbit IgG, HRP Linked Antibody #7074  
TMB Solution #7004  
Stop Solution #7002

**\* NOTE:** Use of an automated microplate washer as well as centrifugation of plates when appropriate, greatly improves reproducibility.

Please contact Cell Signaling Technology for HTS-ready antibodies (PBS formulated and carrier-free), and detailed peptide substrate sequence information.  
Email: drugdiscovery@cellsignal.com