

Store at
-20C
#15125**BRD7 (D9K2T) Rabbit mAb**

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W, IHC-P	H	Endogenous	85	Rabbit IgG	#Q9NPI1	29117

Product Usage Information**Application**

Western Blotting
Immunohistochemistry (Paraffin)

Dilution

1:1000
1:800

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

BRD7 (D9K2T) recognizes endogenous levels of total BRD7 protein.

Source / Purification

Monoclonal antibody is produced by immunizing animals with recombinant protein specific to the carboxy terminus of human BRD7 protein.

Background

Bromodomain-containing protein 7 (BRD7, BP75, CELTIX-1) is a conserved bromodomain-containing protein that was first identified in a screen for proteins that interact with the PDZ domain of PSD95 (1). Subsequent studies identified BRD7 as a major component of SWI/SNF chromatin remodeling complexes, where it was shown to interact directly with acetylated histones to regulate gene transcription (2,3). BRD7 also interacts with p53, and was shown to participate directly in p53-dependent transcriptional regulation (4). Loss-of-function BRD7 mutations were identified in a subset of wild-type p53 breast cancer tumor samples, implicating BRD7 as a putative tumor-suppressor of potential clinical significance (5). BRD7 also associates with the BRCA1 protein, an interaction that facilitates recruitment of BRCA1 to the ERα gene promoter (6).

Background References

- Cuppen, E. et al. (1999) *FEBS Lett* 459, 291-8.
- Peng, C. et al. (2006) *J Cell Biochem* 97, 882-92.
- Kaesler, M.D. et al. (2008) *J Biol Chem* 283, 32254-63.
- Burrows, A.E. et al. (2010) *Proc Natl Acad Sci U S A* 107, 14280-5.
- Drost, J. et al. (2010) *Nat Cell Biol* 12, 380-9.
- Harte, M.T. et al. (2010) *Cancer Res* 70, 2538-47.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IHC-P:** Immunohistochemistry (Paraffin)

Cross-Reactivity Key

H: Human

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

SignalStain is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for

Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.

Orders: 877-616-CELL (2355) • orders@cellsignal.com • Support: 877-678-TECH (8324) • info@cellsignal.com • Web: cellsignal.com
For Research Use Only. Not for Use in Diagnostic Procedures.