

Store at
4°C

CUT&RUN Antibody Binding Buffer



Cell Signaling
TECHNOLOGY®

#15338

2 x 2.5 mL

Support: +1-978-867-2388 (U.S.)
www.cellsignal.com/support

Orders: 877-616-2355 (U.S.)
orders@cellsignal.com

New 01/21

For Research Use Only. Not For Use In Diagnostic Procedures.

Description: The CUT&RUN Antibody Binding Buffer provides enough reagent to support 48 CUT&RUN assays. This product is formulated for optimal performance in the CUT&RUN assay and each lot is tested and validated using the CUT&RUN Assay Kit #86652. An appropriate amount of Digitonin Solution #16359, 100X Spermidine #27287, and Protease Inhibitor Cocktail (200X) #7012 should be added to this product right before use. Please keep on ice during use.

Background: Like the chromatin immunoprecipitation (ChIP) assay, Cleavage Under Targets and Release Using Nuclease (CUT&RUN) is a powerful and versatile technique used for probing protein-DNA interactions within the natural chromatin context of the cell (1-4). CUT&RUN provides a rapid, robust, and true low cell number assay for detection of protein-DNA interactions in the cell. Unlike the ChIP assay, CUT&RUN is free from formaldehyde cross-linking, chromatin fragmentation, and immunoprecipitation, making it a much faster and more efficient method for enriching protein-DNA interactions and identifying target genes. CUT&RUN can be performed in less than one day, from live cells to purified DNA, and has been shown to work with as few as 500-1,000 cells per assay (1,2). Instead of fragmenting all of the cellular chromatin as done in ChIP, CUT&RUN utilizes an antibody-targeted digestion of chromatin, resulting in much lower background signal than seen in the ChIP assay. As a result, CUT&RUN requires only 1/10th the sequencing depth that is required for ChIP-seq assays (1,2). Finally, the inclusion of simple spike-in control DNA allows for accurate quantification and normalization of target-protein binding that is not possible with the ChIP method. This provides for effective normalization of signal between samples and between experiments.

Storage: Store CUT&RUN Antibody Binding Buffer at 4°C. This product is stable for at least 12 months.

Please visit www.cellsignal.com for a complete listing of recommended companion products.

Directions for Use: For the CUT&RUN assay, we recommend adding 1 µl 100X Spermidine #27287, 0.5 µl Protease Inhibitor Cocktail (200X) #7012, and 2.5 µl Digitonin Solution #16359 to 96 µl CUT&RUN Antibody Binding Buffer (100 µl per reaction) right before use. Please keep on ice during use. The volume of Digitonin Solution #16359 in the CUT&RUN Antibody Binding Buffer can be adjusted based on the specific cell line.

Background References:

- (1) Skene, P.J. and Henikoff, S. (2017) *Elife* 6, pii: e21856. doi: 10.7554/eLife.21856.
- (2) Skene, P.J. et al. (2018) *Nat Protoc* 13, 1006-19.
- (3) Meers, M.P. et al. (2019) *Elife* 8, pii: e46314. doi: 10.7554/eLife.46314.
- (4) Meers, M.P. et al. (2019) *Mol Cell* 75, 562-575.e5.

Thank you for your recent purchase. If you would like to provide a review visit cellsignal.com/comments.

www.cellsignal.com

© 2021 Cell Signaling Technology, Inc.

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

Applications: W—Western IP—Immunoprecipitation IHC—Immunohistochemistry ChIP—Chromatin Immunoprecipitation IF—Immunofluorescence F—Flow cytometry E-P—ELISA-Peptide **Species Cross-Reactivity:** H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken Dm—D. melanogaster X—Xenopus Z—zebrafish B—bovine Dg—dog Pg—pig Sc—S. cerevisiae Ce—C. elegans Hr—Horse All—all species expected Species enclosed in parentheses are predicted to react based on 100% homology.