

Store at -20°C

#1592

# Phospho-IRS-1 (Ser1101) Biotinylated Peptide

1.25 ml at 6 µM



Cell Signaling  
TECHNOLOGY®

**Orders** ■ 877-616-CELL (2355)  
orders@cellsignal.com

**Support** ■ 877-678-TECH (8324)  
info@cellsignal.com

**Web** ■ www.cellsignal.com

New 01/08

This product is for *in vitro* research use only and is not intended for use in humans or animals.

**Description:** This biotinylated peptide contains the residues surrounding Ser1101 of IRS-1. The serine residue in the peptide has been chemically phosphorylated in the course of peptide synthesis. This phosphopeptide was generated for use as a positive control for kinase assays using RSK1 Kinase #7398 and RSK2 Kinase #7404, but it may also serve as a positive control in other heterogeneous or homogeneous kinase assays.

**Peptide Core Sequence:** RHSS\*ETF

**Molecular Weight:** 2012 daltons

**Quality Control:** The quality of the biotinylated peptide was evaluated by reverse-phase HPLC and by mass spectrometry.

**Directions for Use:** This phosphorylated peptide can be detected with the Phospho-IRS-1 (Ser1101) Antibody #2385. A sample kinase assay protocol is attached.

**Entrez-Gene ID** # 3667

**Swiss-Prot Acc.** # P35568

**Storage:** Supplied in 0.0001% DMSO. Store at -20°C.

**Companion Products:**

RSK1 Kinase #7398

RSK2 Kinase #7404

Phospho-IRS-1 (Ser1101) Antibody #2385

IRS-1 (Ser1101) Biotinylated Peptide #1046

## Protocol for Serine/Threonine Kinase Assay

**IMPORTANT:** Use of an automated microplate washer as well as centrifugation of plates when appropriate, greatly improves reproducibility.

### A Additional Solutions and Reagents (Not included)

1. **Wash Buffer:** 1X PBS, 0.05% Tween-20 (PBS/T)
2. Bovine Serum Albumin (BSA)
3. **Stop Buffer:** 50 mM EDTA pH 8
4. Kinase Buffer (10X) #9802
5. ATP (10 mM) #9804
6. Active kinase (See companion products)
7. Primary antibody (See companion products)

### B Suggested Protocol for 100 Assays

1. Add 100  $\mu$ l 10 mM ATP to 1.25 ml 6-12  $\mu$ M substrate peptide. Adjust the mixture with  $dH_2O$  to 2.5 ml to make 2X ATP/substrate cocktail ([ATP]=400  $\mu$ M, [substrate] = 3-6  $\mu$ M).
2. Transfer enzyme from  $-80^{\circ}C$  to ice. Allow enzyme to thaw on ice.
3. **Microcentrifuge briefly at  $4^{\circ}C$  to bring liquid to the bottom of the vial. Return immediately to ice.**
4. Add 1 ml 10X kinase buffer [250 mM Tris-HCl pH 7.5, 100 mM  $MgCl_2$ , 1 mM  $Na_3VO_4$ , 50 mM  $\beta$ -glycerophosphate, 20 mM dithiothreitol (DTT)] to 1.5 ml  $dH_2O$  to make 2.5 ml 4X reaction buffer.
5. Dilute enzyme in 1.25 ml of 4X reaction buffer to make 4X reaction cocktail ([enzyme]=0.8-8.0 ng/ $\mu$ l in 4X reaction cocktail).
6. Add 12.5  $\mu$ l of the 4X reaction cocktail to 12.5  $\mu$ l/well of prediluted compound of interest (usually around 10  $\mu$ M) and incubate for 5 minutes at room temperature.
7. Add 25  $\mu$ l of 2X ATP/substrate cocktail to 25  $\mu$ l/well preincubated reaction cocktail/compound.

#### Final Assay Conditions for a 50 $\mu$ l Reaction

25 mM Tris-HCl (pH7.5)  
10 mM  $MgCl_2$   
5 mM  $\beta$ -glycerophosphate  
0.1 mM  $Na_3VO_4$   
2 mM DTT  
200  $\mu$ M ATP  
1.5-3  $\mu$ M peptide  
10-100 ng kinase

8. Incubate reaction plate at room temperature for 30 minutes.
9. Add 50  $\mu$ l/well Stop Buffer (50 mM EDTA, pH 8) to stop the reaction.
10. Transfer 25  $\mu$ l of each reaction to a 96-well streptavidin-coated plate containing 75  $\mu$ l  $dH_2O$ /well and incubate at room temperature for 60 minutes.
11. Wash three times with 200  $\mu$ l/well PBS/T.
12. Dilute primary antibody in PBS/T with 1% BSA. \*Add 100  $\mu$ l/well primary antibody.(1:500 dilution for mouse mAb or 1:1000 dilution for rabbit mAb or polyclonal antibody)
13. Incubate at  $37^{\circ}C$  for 120 minutes.
14. Wash three times with 200  $\mu$ l/well PBS/T.
15. For DELFIA<sup>®</sup> or Colorimetric ELISA detection methods please use the following protocols.

DELFIA<sup>®</sup> is a registered trademark of PerkinElmer Life Sciences

### DELFIA<sup>®</sup> Assay

1. Prepare appropriate dilution of Europium labeled secondary antibody in PBS/T with 1% BSA (1:500 dilution for anti-mouse IgG or 1:1000 for anti-rabbit IgG).
2. Add 100  $\mu$ l/well secondary antibody solution.
3. Incubate at room temperature for 30 minutes.
4. \*Wash five times with 200  $\mu$ l/well PBS/T.
5. Add 100  $\mu$ l/well DELFIA<sup>®</sup> Enhancement Solution.
6. Incubate at room temperature for 5 minutes.
7. Read plate using a Time Resolved Fluorescent plate reader using the following settings;
  - a. Excitation Filter: 340 nm
  - b. Emission Filter: 615 nm
  - c. Delay\*\*: 400  $\mu$ s
 \*\* Delay time is the delay from the excitation pulse to the beginning of the measurement.

### Companion Products for DELFIA<sup>®</sup>

DELFIA<sup>®</sup> Europium-labeled Anti-mouse IgG (PerkinElmer Life Sciences #AD0124)  
DELFIA<sup>®</sup> Europium-labeled Anti-rabbit IgG (PerkinElmer Life Sciences #AD0105)  
DELFIA<sup>®</sup> Enhancement Solution (PerkinElmer Life Sciences #1244-105)  
DELFIA<sup>®</sup> Streptavidin coated, 96-well, yellow plate (PerkinElmer Life Sciences AAAND-0005)

### Colorimetric ELISA Assay

1. Prepare appropriate dilution of HRP labeled secondary antibody in PBS/T with 1% BSA (1:500 dilution for anti-mouse IgG or 1:1000 for anti-rabbit IgG).
2. Add 100  $\mu$ l/well secondary antibody solution.
3. Incubate at room temperature for 30 minutes.
4. \*Wash five times with 200  $\mu$ l/well PBS/T.
5. Add 100  $\mu$ l/well TMB substrate.
6. Incubate at room temperature for 15 minutes.
7. Add 100  $\mu$ l/well of stop solution.
8. Mix well.
9. Read the absorbance at 450 nm with a microtiter plate reader.

### Companion Products For Colorimetric ELISA Assay

Anti-mouse IgG, HRP Linked Antibody #7076  
Anti-rabbit IgG, HRP Linked Antibody #7074  
TMB Solution #7004  
Stop Solution #7002

\* **NOTE:** Use of an automated microplate washer as well as centrifugation of plates when appropriate, greatly improves reproducibility.

Please contact Cell Signaling Technology for HTS-ready antibodies (PBS formulated and carrier-free), and detailed peptide substrate sequence information.  
Email: drugdiscovery@cellsignal.com