Store at -20C	Phospho-Akt1 (Ser473) Mat Pair			Antibody	C T	Cell Signaling	
					Orders:	877-616-CELL (2355) orders@cellsignal.com	
55					Support:	877-678-TECH (8324)	
60	Species Cross Reactivity: H M R Mk	UniProt ID: #P31749	Entrez-Gene Id: #207		Web:	info@cellsignal.com cellsignal.com	
#1				3 Trask Lane	Danvers Mas	sachusetts 01923 USA	

For Research Use Only. Not for Use in Diagnostic Procedures.

Product Includes		Product #	Quantity	Isotype/Source	
Akt1 (C73H10) Rabbit mAb (BSA and	84505	100 µg	Rabbit IgG		
Phospho-Akt (Ser473) (193H12) Rabl	92982	100 µg	Rabbit IgG		
Description	ion The Phospho-Akt1 (Ser473) Matched Antibody Pair is ideal for use with immunoassay technologies and high-throughput ELISA platforms requiring antibody pairs with specialized or custom antibody labeling. Labels include fluorophores, lanthanides, biotin, and beads. Platforms requiring conjugated Matched Antibody Pairs include MSD, Quanterix Simoa, Alpha Technology (AlphaScreen, AlphaLISA, LANCE, HTRF), and Luminex.				
	antibody-pairs.	our projects forward	, laster at cst-so	lience.com/matched-	
Specificity/Sensitivity	This kit detects proteins from the indicated also detect homologous proteins from othe	ned through in-	house testing, but may		
Storage	Store at -20°C. <i>This product will freeze at -20°C so it is recommended to aliquot into single-use vials to avoid multiple freeze/thaw cycles</i> . A slight precipitate may be present and can be dissolved by gently vortexing. This will not interfere with antibody performance.				
Directions for Use	Matched Antibody Pairs include capture and detection antibodies to non-overlapping epitopes. Optimal dilutions/concentrations should be determined by the end user.				
Formulation	Supplied in 1X PBS (10 mM Na $_2$ HPO $_4$, 3 mM KCl, 2 mM KH $_2$ PO $_4$, and 140 mM NaCl (pH 7.8)). BSA and Azide Free.				
Background	Akt, also referred to as PKB or Rac, plays a critical role in controlling cell survival and apoptosis (1-3). This protein kinase is activated by insulin and various growth and survival factors to function in a wortmannin-sensitive pathway involving PI3 kinase (2,3). Akt is activated by phospholipid binding and activation loop phosphorylation at Thr308 by PDK1 (4) and by phosphorylation within the carboxy terminus at Ser473. The previously elusive PDK2 responsible for phosphorylation of Akt at Ser473 has been identified as mammalian target of rapamycin (mTOR) in a rapamycin-insensitive complex with rictor and Sin1 (5,6). Akt promotes cell survival by inhibiting apoptosis through phosphorylation and inactivation of several targets, including Bad (7), forkhead transcription factors (8), c-Raf (9), and caspase-9. PTEN phosphatase is a major negative regulator of the PI3K/Akt signaling pathway (10). LY294002 is a specific PI3 kinase inhibitor (11). Another essential Akt function is the regulation of glycogen synthesis through phosphorylation and nactivation of GSK-30 and β (12,13). Akt may also play a role in insulin stimulation of glucose transport (12). In addition to its role in survival and glycogen synthesis, Akt is involved in cell cycle regulation by preventing GSK-3β-mediated phosphorylation and degradation of cyclin D1 (14) and by negatively regulating the cyclin-dependent kinase inhibitors p27 Kip1 (15) and p21 Waf1/Cip1 (16). Akt also plays a critical role in cell growth by directly phosphorylating mTOR in a rapamycin-sensitive complex containing raptor (17). More importantly, Akt phosphorylates and inactivates tuberin (TSC2), an inhibitor of mTOR within the mTOR-raptor complex (18,19).				
Background References	 Franke, T.F. et al. (1997) <i>Cell</i> 88, 435-7. Burgering, B.M. and Coffer, P.J. (1995) <i>Na</i> Franke, T.F. et al. (1995) <i>Cell</i> 81, 727-36. Alessi, D.R. et al. (1996) <i>EMBO J</i> 15, 6541- Sarbassov, D.D. et al. (2005) <i>Science</i> 307, Jacinto, E. et al. (2006) <i>Cell</i> 127, 125-37. Cardone, M.H. et al. (1998) <i>Science</i> 282, 1 Brunet, A. et al. (1999) <i>Cell</i> 96, 857-68. Zimmermann, S. and Moelling, K. (1999) Cantley, L.C. and Neel, B.G. (1999) <i>Proc</i>. Vlahos, C.J. et al. (2001) <i>FEBS Lett</i> 492, 1 	<i>ture</i> 376, 599-602. 51. 1098-101. 1318-21. <i>Science</i> 286, 1741-4. <i>Natl Acad Sci USA</i> 96, , 5241-8. 199-203.	4240-5.		

	 13. Cross, D.A. et al. (1995) <i>Nature</i> 378, 785-9. 14. Diehl, J.A. et al. (1998) <i>Genes Dev</i> 12, 3499-511. 15. Gesbert, F. et al. (2000) <i>J Biol Chem</i> 275, 39223-30. 16. Zhou, B.P. et al. (2001) <i>Nat Cell Biol</i> 3, 245-52. 17. Navé, B.T. et al. (1999) <i>Biochem J</i> 344 Pt 2, 427-31. 18. Inoki, K. et al. (2002) <i>Nat Cell Biol</i> 4, 648-57. 19. Manning, B.D. et al. (2002) <i>Mol Cell</i> 10, 151-62.
Trademarks and Patents	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
	U.S. Patent No. 7,429,487, foreign equivalents, and child patents deriving therefrom.
	All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.
Limited Uses	Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.
	Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.

Orders: 877-616-CELL (2355) • orders@cellsignal.com • Support: 877-678-TECH (8324) • info@cellsignal.com • Web: cellsignal.com For Research Use Only. Not for Use in Diagnostic Procedures.