

TKS5 Antibody



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, IP	Reactivity: H	Sensitivity: Endogenous	MW (kDa): 150	Source/Isotype: Rabbit	UniProt ID: #Q5TCZ1	Entrez-Gene Id: 9644
-------------------------------	-------------------------	-----------------------------------	-------------------------	----------------------------------	-------------------------------	--------------------------------

Product Usage Information

Application

Western Blotting
Immunoprecipitation

Dilution

1:1000
1:50

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

TKS5 Antibody recognizes endogenous levels of total TKS5 protein. This antibody cross-reacts with a 300 kDa protein of unknown identity.

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Asp910 of human TKS5 protein. Antibodies are purified by protein A and peptide affinity chromatography.

Background

TKS5 (SH3PXD2A, FISH) is a scaffold protein expressed on invadosomes of both normal and transformed cell lines. Research studies suggest that TKS5 is functionally required for both the formation and invasive behavior of invadosomes (1, 2). TKS5 has an N-terminal PX domain that mediates invadosome initiation and localization of MMP-rich vesicles to the invadosome (3). TKS5 also has five SH3 domains, which recruit ADAM family proteinases to the invadosome to degrade extracellular matrix. These SH3 domains interact with adaptor proteins to facilitate F-actin polymerization during invadosome formation (4-6). Src tyrosine kinase has been shown to phosphorylate TKS5 at Tyr557 and Tyr619, which was shown to be necessary and sufficient for TKS5-mediated invadopodia formation and invasion (7). Elevated TKS5 expression is positively associated with invasive behavior of cancer cells, suggesting TKS5 may have prognostic potential in cancer (8, 9).

Background References

1. Paterson, E.K. and Courtneidge, S.A. (2017) *FEBS J* doi:10.1111/febs.14123.
2. Courtneidge, S.A. (2012) *Biochem Soc Trans* 40, 129-32.
3. Jacob, A. et al. (2016) *J Cell Sci* 129, 4341-4353.
4. Abram, C.L. et al. (2003) *J Biol Chem* 278, 16844-51.
5. Crimaldi, L. et al. (2009) *Exp Cell Res* 315, 2581-92.
6. Stylli, S.S. et al. (2009) *J Cell Sci* 122, 2727-40.
7. Burger, K.L. et al. (2014) *Prostate* 74, 134-48.
8. Stylli, S.S. et al. (2014) *Oncol Rep* 32, 989-1002.
9. Blouw, B. et al. (2015) *PLoS One* 10, e0121003.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IP:** Immunoprecipitation

Cross-Reactivity Key

H: Human

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless

separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.