



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Store at -20C
#2052

PKD/PKC μ Antibody

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, FC-FP	Reactivity: H M Mk	Sensitivity: Endogenous	MW (kDa): 115	Source/Isotype: Rabbit	UniProt ID: #Q15139	Entrez-Gene Id: 5587
----------------------------------	------------------------------	-----------------------------------	-------------------------	----------------------------------	-------------------------------	--------------------------------

Product Usage Information

Application

Western Blotting
Flow Cytometry (Fixed/Permeabilized)

Dilution

1:1000
1:50

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

PKD/PKC μ Antibody detects endogenous levels of total PKD1/PKC μ protein. This antibody may also react with isoform PKD2, in some species.

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Ser916 of mouse PKD. Antibodies are purified by protein A and peptide affinity chromatography.

Background

Activation of PKC is one of the earliest events in a cascade leading to a variety of cellular responses, such as secretion, gene expression, proliferation, and muscle contraction (1,2). Protein kinase D (PKD), also called PKC μ , is a serine/threonine kinase whose activation is dependent on the phosphorylation of two activation loop sites, Ser744 and Ser748, via a PKC-dependent signaling pathway (3-5). In addition to the two activation loop sites, the carboxy-terminal Ser916 has been identified as an autophosphorylation site for PKD/PKC μ . Phosphorylation at Ser916 correlates with PKD/PKC μ catalytic activity (6).

Background References

1. Nishizuka, Y. (1984) *Nature* 308, 693-698.
2. Keranen, L.M. (1995) *Curr. Biol.* 5, 1394-1403.
3. Valverde, A.M. et al. (1994) *Proc. Natl. Acad. Sci.* 91, 8572-8576.
4. Johannes, F.J. et al. (1994) *J. Biol. Chem.* 269, 6140-6148.
5. Iglesias, T. et al. (1998) *J. Biol. Chem.* 273, 27662-27667.
6. Matthews, S.A. et al. (1999) *J. Biol. Chem.* 274, 26543-26549.
7. Wood, C.D. et al. (2005) *J Biol Chem* 280, 6245-51.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween@ 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **FC-FP:** Flow Cytometry (Fixed/Permeabilized)

Cross-Reactivity Key

H: Human **M:** Mouse **Mk:** Monkey

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in

any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.

Orders: 877-616-CELL (2355) • orders@cellsignal.com • Support: 877-678-TECH (8324) • info@cellsignal.com • Web: cellsignal.com
For Research Use Only. Not for Use in Diagnostic Procedures.