



**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Store at -20C  
#2088

## LEDGF (C57G11) Rabbit mAb

**For Research Use Only. Not for Use in Diagnostic Procedures.**

<b>Applications:</b> W, IHC-P, IF-IC, FC-FP	<b>Reactivity:</b> H M R	<b>Sensitivity:</b> Endogenous	<b>MW (kDa):</b> 75	<b>Source/Isotype:</b> Rabbit IgG	<b>UniProt ID:</b> #O75475	<b>Entrez-Gene Id:</b> 11168
------------------------------------------------	-----------------------------	-----------------------------------	------------------------	--------------------------------------	-------------------------------	---------------------------------

### Product Usage Information

#### Application

Western Blotting  
Immunohistochemistry (Paraffin)  
Immunofluorescence (Immunocytochemistry)  
Flow Cytometry (Fixed/Permeabilized)

#### Dilution

1:1000  
1:50  
1:400  
1:100

### Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

For a carrier free (BSA and azide free) version of this product see product #96208.

### Specificity/Sensitivity

LEDGF (C57G11) Rabbit mAb detects endogenous levels of LEDGF protein.

### Species predicted to react based on 100% sequence homology

Monkey

### Source / Purification

Monoclonal antibody is produced by immunizing animals with synthetic peptide corresponding to amino acids surrounding Lys461 of human LEDGF.

### Background

The LEDGF (lens epithelium-derived growth factor) gene encodes splice variants P75 and P52 (1). LEDGF (C57G11) Rabbit mAb detects the 75 kDa LEDGF splice form. Both LEDGF variants have transcriptional coactivator functions that may regulate gene transcription and mRNA splicing of a variety of proteins that are essential for cell growth and survival. (2,3). LEDGF has also been identified as a factor that can, in serum free media, stimulate the growth and survival of lens epithelial cells and other cell types such as retinal photoreceptor cells, COS7 cells, skin fibroblasts, and keratinocytes (2). In addition to its function as a transcriptional coactivator, LEDGF has been shown to be involved in the pathogenesis of atopic dermatitis (4) and in the activity of HIV1 integrase. LEDGF binds chromatin throughout the cell cycle, and may act as a chromatin docking factor for HIV integrase and other lentiviral pre-integration complexes. It protects HIV integrase from degradation, and influences the pattern of viral integration into the cell genome (5).

### Background References

1. Ge, H. et al. (1998) *EMBO J* 17, 6723-9.
2. Singh, D.P. et al. (2000) *Biochem Biophys Res Commun* 267, 373-81.
3. Singh, D.P. et al. (2001) *Biochem Biophys Res Commun* 283, 943-55.
4. Sugiura, K. et al. (2007) *J Invest Dermatol* 127, 75-80.
5. Poeschla, E.M. (2008) *Cell Mol Life Sci* 65, 1403-24.

### Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

### Western Blot Buffer

**IMPORTANT:** For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween@ 20 at 4°C with gentle shaking, overnight.

### Applications Key

**W:** Western Blotting **IHC-P:** Immunohistochemistry (Paraffin) **IF-IC:** Immunofluorescence (Immunocytochemistry) **FC-FP:** Flow Cytometry (Fixed/Permeabilized)

### Cross-Reactivity Key

**H:** Human **M:** Mouse **R:** Rat

### Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

U.S. Patent No. 7,429,487, foreign equivalents, and child patents deriving therefrom.

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

## Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.