## **P-Cadherin Antibody**



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

## For Research Use Only. Not for Use in Diagnostic Procedures.

<b>Applications:</b> W, IF-IC	Reactivity: H	<b>Sensitivity:</b> Endogenous	<b>MW (kDa):</b> 120	Source/Isotype: Rabbit	UniProt ID: #P22223	Entrez-Gene Id: 1001
Product Usage Information		<b>Application</b> Western Blotting Immunofluorescence	(Immunocytochen	nistry)		<b>Dilution</b> 1:1000 1:100
Storage		Supplied in 10 mM so 20°C. Do not aliquot t		5), 150 mM NaCl, 100 μg,	/ml BSA and 50% g	lycerol. Store at –
Specificity/Sensitivity		P-Cadherin Antibody recognizes endogenous levels of total P-cadherin protein. The antibody does not cross-react with other cadherin family members.				
Source / Purification		Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to human P-cadherin. Antibodies are purified using protein A and peptide affinity chromatography.				
Background		Cadherins are a superfamily of transmembrane glycoproteins that contain cadherin repeats of approximately 100 residues in their extracellular domain. Cadherins mediate calcium-dependent cell-cell adhesion and play critical roles in normal tissue development (1). The classic cadherin subfamily includes N-, P-, R-, B-, and E-cadherins, as well as about ten other members that are found in adherens junctions, a cellular structure near the apical surface of polarized epithelial cells. The cytoplasmic domain of classical cadherins interacts with $\beta$ -catenin, $\gamma$ -catenin (also called plakoglobin), and p120 catenin. $\beta$ -catenin and $\gamma$ -catenin associate with $\alpha$ -catenin, which links the cadherin-catenin complex to the actin cytoskeleton (1,2). While $\beta$ - and $\gamma$ -catenin play structural roles in the junctional complex, p120 regulates cadherin adhesive activity and trafficking (1-4). Investigators consider E-cadherin an active suppressor of invasion and growth of many epithelial cancers (1-3). Research studies indicate that cancer cells have upregulated N-cadherin in addition to loss of E-cadherin. This change in cadherin expression is called the "cadherin switch." N-cadherin cooperates with the FGF receptor, leading to overexpression of MMP-9 and cellular invasion (3). Research studies have shown that in endothelial cells, VE-cadherin signaling, expression, and localization correlate with vascular permeability and tumor angiogenesis (5,6). Investigators have also demonstrated that expression of P-cadherin, which is normally present in epithelial cells, is also altered in ovarian and other human cancers (7,8).				
Background References		<ol> <li>Wheelock, M.J. and Johnson, K.R. (2003) <i>Annu Rev Cell Dev Biol</i> 19, 207-35.</li> <li>Christofori, G. (2003) <i>EMBO J</i> 22, 2318-23.</li> <li>Hazan, R.B. et al. (2004) <i>Ann N Y Acad Sci</i> 1014, 155-63.</li> <li>Bryant, D.M. and Stow, J.L. (2004) <i>Trends Cell Biol</i> 14, 427-34.</li> <li>Rabascio, C. et al. (2004) <i>Cancer Res</i> 64, 4373-7.</li> <li>Yamaoka-Tojo, M. et al. (2006) <i>Arterioscler Thromb Vasc Biol</i> 26, 1991-7.</li> <li>Patel, I.S. et al. (2003) <i>Int J Cancer</i> 106, 172-7.</li> <li>Sanders, D.S. et al. (2000) <i>J Pathol</i> 190, 526-30.</li> </ol>				
Species Reactivit	у	Species reactivity is de	etermined by testin	g in at least one approve	ed application (e.g.,	western blot).
Western Blot Buf	ffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.				

**Applications Key** 

W: Western Blotting IF-IC: Immunofluorescence (Immunocytochemistry)

**Cross-Reactivity Key** 

H: Human

**Trademarks and Patents** 

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

Alexa Fluor is a registered trademark of Life Technologies Corporation.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

## **Limited Uses**

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party. whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.