



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Store at -20C
#2296

GM130 Antibody

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, IP	Reactivity: H	Sensitivity: Endogenous	MW (kDa): 130	Source/Isotype: Rabbit	UniProt ID: #Q08379	Entrez-Gene Id: 2801
-------------------------------	-------------------------	-----------------------------------	-------------------------	----------------------------------	-------------------------------	--------------------------------

Product Usage Information

Application

Western Blotting
Immunoprecipitation

Dilution

1:1000
1:25

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

GM130 Antibody detects endogenous levels of total GM130 protein.

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to the amino-terminal sequence of human GM130.

Background

The Golgi apparatus functions in the modification, organization, and transport of proteins and membranes targeted to other parts of the cell, such as the plasma membrane, lysosomes, and endosomes. This regulated transport is important for appropriate protein localization, secretion, and signal transduction. Members of the Golgin family of proteins, including GM130, Giantin, p115, and GRASP65, are defined by their presence in the Golgi matrix and by their long coiled-coil domains. Golgin function, which is regulated in part by small GTPases of the Rab and Arl families, includes establishing and maintaining Golgi structure and transport (reviewed in 1). The Golgi cisternae are stacked and linked laterally to form a ribbon. GRASP65 and GM130 are required for membrane fusion events that mediate ribbon formation during Golgi assembly. These lateral fusion events allow for uniform distribution of Golgi enzymes (2). GM130 and Giantin interact with the transport factor p115 to facilitate endoplasmic reticulum (ER)-Golgi transport (3). GM130 is also involved in the transport of the ether-a-go-go-related (hERG) potassium ion channel. Inappropriate hERG localization may be an underlying cause of long QT syndrome, a hereditary and potentially fatal cardiac arrhythmia (4). Further, GM130 was implicated in signal transduction regulating invasion, migration, and cell polarization via its interaction with and activation of serine/threonine kinases YSK1 and Mst4 (5).

Background References

1. Barr, F.A. and Short, B. (2003) *Curr. Opin. Cell Biol.* 15, 405-413.
2. Puthenveedu, M.A. et al. (2006) *Nat. Cell Biol.* 8, 238-248.
3. Alvarez, C. et al. (2001) *J. Biol. Chem.* 276, 2693-2700.
4. Roti, E.C. et al. (2002) *J. Biol. Chem.* 277, 47779-47785.
5. Preisinger, C. et al. (2004) *J. Cell Biol.* 164, 1009-1020.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween@ 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IP:** Immunoprecipitation

Cross-Reactivity Key

H: Human

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no

force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.