



**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Store at -20C  
#2391

## Wnt3a Antibody

**For Research Use Only. Not for Use in Diagnostic Procedures.**

<b>Applications:</b> W	<b>Reactivity:</b> M	<b>Sensitivity:</b> Transfected Only	<b>MW (kDa):</b> 42	<b>Source/Isotype:</b> Rabbit	<b>UniProt ID:</b> #P56704	<b>Entrez-Gene Id:</b> 89780
---------------------------	-------------------------	---	------------------------	----------------------------------	-------------------------------	---------------------------------

<b>Product Usage Information</b>	<b>Application</b> Western Blotting	<b>Dilution</b> 1:1000
<b>Storage</b>	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.	
<b>Specificity/Sensitivity</b>	Wnt3a antibody detects transfected Wnt3a protein in L cells and the conditioned medium from L/Wnt3a cells. Based on homology, it is expected to also recognize Wnt3.	
<b>Species predicted to react based on 100% sequence homology</b>	Human	
<b>Source / Purification</b>	Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Asp294 of human Wnt3a. Antibodies are purified by protein A and peptide affinity chromatography.	
<b>Background</b>	<p>The Wnt family includes several secreted glycoproteins that play important roles in animal development (1). There are 19 Wnt genes in the human genome that encode functionally distinct Wnt proteins (2). Wnt members bind to the Frizzled family of seven-pass transmembrane proteins and activate several signaling pathways (3). The canonical Wnt/β-catenin pathway also requires a coreceptor from the low-density lipoprotein receptor family (4). Aberrant activation of Wnt signaling pathways is involved in several types of cancers (5).</p> <p>Wnt3a protein is palmitoylated and can function as a growth factor for hematopoietic stem cells (6). Although functionally distinct, Wnt3a shows high homology to Wnt3 (7).</p>	
<b>Background References</b>	<ol style="list-style-type: none"> <li>1. Cadigan, K.M. and Nusse, R. (1997) <i>Genes Dev.</i> 11, 3286-3305.</li> <li>2. Moon, R.T. et al. (1997) <i>Trends Genet.</i> 13, 157-162.</li> <li>3. Kohn, A.D. and Moon, R.T. (2004) <i>Cell Calcium</i> 38, 439-446.</li> <li>4. Logan, C.Y. and Nusse, R. (2004) <i>Annu. Rev. Cell Dev. Biol.</i> 20, 781-810.</li> <li>5. Giles, R.H. et al. (2003) <i>Biochim. Biophys. Acta.</i> 1653, 1-24.</li> <li>6. Willert, K. et al. (2003) <i>Nature</i> 423, 448-52.</li> <li>7. Katoh, M. (2001) <i>Int. J. Oncol.</i> 19, 977-82.</li> </ol>	
<b>Species Reactivity</b>	Species reactivity is determined by testing in at least one approved application (e.g., western blot).	
<b>Western Blot Buffer</b>	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.	
<b>Applications Key</b>	<b>W:</b> Western Blotting	
<b>Cross-Reactivity Key</b>	<b>M:</b> Mouse	
<b>Trademarks and Patents</b>	<p>Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.</p> <p>All other trademarks are the property of their respective owners. Visit <a href="http://cellsignal.com/trademarks">cellsignal.com/trademarks</a> for more information.</p>	
<b>Limited Uses</b>	<p>Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.</p>	

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.