

**NOP2 (G754) Antibody**

**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

**For Research Use Only. Not for Use in Diagnostic Procedures.**

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W, IP	H	Endogenous	110	Rabbit	#P46087	4839

**Product Usage Information****Application**

Western Blotting  
Immunoprecipitation

**Dilution**

1:1000  
1:50

**Storage**

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. *Do not aliquot the antibody.*

**Specificity/Sensitivity**

NOP2 (G754) Antibody recognizes endogenous levels of total NOP2 protein.

**Source / Purification**

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Gly754 of human NOP2 protein. Antibodies are purified by peptide affinity chromatography.

**Background**

Chemical modifications of RNA regulate many cellular processes. One particular RNA modification, 5-methylcytosine (5-mC), regulates ribosome assembly, translation, and RNA stability (1). In eukaryotes, this modification is added to RNA by the DNA methyltransferase homologue 2 protein (DNMT2; also known as TRDMT1), and also by members of the NOL1/NOP2/SUN domain (NSUN) family of proteins. NSUN proteins are putative S-adenosylmethionine (SAM)-dependent methyltransferases that carry out their enzymatic activity by utilizing two cysteine residues in their active sites (2). There are currently seven known members of this family, consisting of NOP2 (NSUN1) and NSUN2-7.

NOP2, also known as NSUN1, is an 89 kDa member of the NSUN protein family that specifically targets and methylates 28S rRNA (3). In humans, methylation of C4413 in the 28S rRNA by NOP2 is thought to increase stability of the ribosome. NOP2 is strongly overexpressed in many cancers, including colorectal and lung cancer, where it may contribute to tumorigenesis by recruiting telomerase to the cyclin D1 promoter and activating gene expression (4-6). NOP2 also interacts with BRD4 and RNA polymerase II, suggesting additional roles for NOP2 in regulating transcription (1,7).

**Background References**

1. Trixl, L. and Lusser, A. (2019) *Wiley Interdiscip Rev RNA* 10, e1510.
2. Bohnsack, K.E. et al. (2019) *Genes (Basel)* 10, pii: E102. doi: 10.3390/genes10020102.
3. Bourgeois, G. et al. (2015) *PLoS One* 10, e0133321.
4. Ueki, T. et al. (1997) *Hum Pathol* 28, 74-9.
5. Uchiyama, B. et al. (1997) *Clin Cancer Res* 3, 1873-7.
6. Hong, J. et al. (2016) *J Cell Sci* 129, 1566-79.
7. Cheng, J.X. et al. (2018) *Nat Commun* 9, 1163.

**Species Reactivity**

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

**Western Blot Buffer**

**IMPORTANT:** For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

**Applications Key**

**W:** Western Blotting **IP:** Immunoprecipitation

**Cross-Reactivity Key**

**H:** Human

**Trademarks and Patents**

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

**Limited Uses**

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's

terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.