

β-Amyloid (1-43) (E8C2D) Rabbit mAb (Alexa Fluor[®] 555 Conjugate)



Orders: 877-616-CELL (2355) orders@cellsignal.com Support: 877-678-TECH (8324) Web: info@cellsignal.com cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: IF-F	Reactivity: H	Sensitivity: Endogenous	Source/Isotype: Rabbit IgG	UniProt ID: #P05067	Entrez-Gene Id: 351
Product Usage Information		Application Immunofluorescence (Fr	ozen)		Dilution 1:50
Storage		Supplied in PBS (pH 7.2), less than 0.1% sodium azide and 2 mg/ml BSA. Store at 4°C. <i>Do not aliquot the antibody. Protect from light. Do not freeze</i> .			
Specificity/Sensitivity		β-Amyloid (1-43) (E8C2D) Rabbit mAb (Alexa Fluor [®] 555 Conjugate) recognizes endogenous levels of total human Aβ-43 protein. This product detects transgenically expressed human APP in mouse models. This antibody weakly cross-reacts with human Aβ-42 protein.			
Species predicte based on 100% s homology	d to react equence	Mouse			
Source / Purification		Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues near the carboxy of human β -Amyloid (1-43) protein.			
Description		This Cell Signaling Technology antibody is conjugated to Alexa Fluor [®] 555 fluorescent dye and tested in-house for direct immunofluorescent analysis in mouse tissue. This antibody is expected to exhibit the same species cross-reactivity as the unconjugated β -Amyloid (1-43) (E8C2D) Rabbit mAb #32098.			
Background		Amyloid β (A β) precursor protein (APP) is a 100-140 kDa transmembrane glycoprotein that exists as several isoforms (1). The amino acid sequence of APP contains the amyloid domain, which can be released by a two-step proteolytic cleavage (1). The extracellular deposition and accumulation of the released A β fragments form the main components of amyloid plaques in Alzheimer's disease (1). APP can be phosphorylated at several sites, which may affect the proteolytic processing and secretion of this protein (2-5). Phosphorylation at Thr668 (a position corresponding to the APP695 isoform) by cyclin-dependent kinase is cell-cycle dependent and peaks during G2/M phase (4). APP phosphorylated at Thr668 exists in adult rat brain and correlates with cultured neuronal differentiation (5,6).			
Background References		1. Selkoe, D.J. (1996) <i>J Biol Chem</i> 271, 18295-8. 2. Caporaso, G.L. et al. (1992) <i>Proc Natl Acad Sci USA</i> 89, 3055-9. 3. Hung, A.Y. and Selkoe, D.J. (1994) <i>EMBO J</i> 13, 534-42. 4. Suzuki, T. et al. (1994) <i>EMBO J</i> 13, 1114-22. 5. Ando, K. et al. (1999) <i>J Neurosci</i> 19, 4421-7. 6. Iijima, K. et al. (2000) <i>J Neurochem</i> 75, 1085-91.			
Species Reactivity		Species reactivity is determined by testing in at least one approved application (e.g., western blot).			
Applications Key		IF-F: Immunofluorescence (Frozen)			
Cross-Reactivity Key		H: Human			
Trademarks and Patents		Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.			
		XP is a registered traden			
		transfer of this product i conducted by the buyer, must not (1) use this pro purposes; (b) testing, an per-test basis; or (c) man	s conditioned on the buy excluding contract resea duct or its components fo alysis or screening servic ufacturing or quality ass	er using the purchas rch or any fee for ser or (a) diagnostic, ther es, or information in urance or quality con	e Technologies Corporation. The ed product solely in research vice research, and the buyer apeutic or prophylactic return for compensation on a trol, and/or (2) sell or transfer e in research. For information on

purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.