

Histone H4 Antibody



Orders ■ 877-616-CELL (2355)
orders@cellsignal.com
Support ■ 877-678-TECH (8324)
info@cellsignal.com
Web ■ www.cellsignal.com

rev. 12/28/15

For Research Use Only. Not For Use In Diagnostic Procedures.


Entrez-Gene ID # 8359
Swiss-Prot Acc. # P62805

Applications	Species Cross-Reactivity*	Molecular Wt.	Source
W, IP Endogenous	H, M, R, Mk, Sc, Dm, Z	11 kDa	Rabbit**


Background: Modulation of chromatin structure plays an important role in the regulation of transcription in eukaryotes. The nucleosome, made up of four core histone proteins (H2A, H2B, H3 and H4), is the primary building block of chromatin (1). The amino-terminal tails of core histones undergo various post-translational modifications, including acetylation, phosphorylation, methylation and ubiquitination (2-5). These modifications occur in response to various stimuli and have a direct effect on the accessibility of chromatin to transcription factors and, therefore, on gene expression (6). In most species, histone H2B is primarily acetylated at lysines 5, 12, 15 and 20 (4,7). Histone H3 is primarily acetylated at Lys9, 14, 18 and 23 (2,3). Acetylation at Lys9 appears to have a dominant role in histone deposition and chromatin assembly in some organisms (2,3). Phosphorylation at Ser10, Ser28 and Thr11 of histone H3 is tightly correlated with chromosome condensation during both mitosis and meiosis (8-10). Phosphorylation of Thr3 of histone H3 is highly conserved among many species and is catalyzed by the kinase haspin. Immunostaining with phospho-specific antibodies in mammalian cells reveals mitotic phosphorylation of H3 Thr3 in prophase and its dephosphorylation during anaphase (11).

Specificity/Sensitivity: Histone H4 Antibody detects endogenous levels of total histone H4 protein. The antibody does not cross-react with other histones.

Source/Purification: Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to the amino-terminal sequence of human histone H4. Antibodies are purified by protein A and peptide affinity chromatography.



Western blot analysis of extracts from various cell types using Histone H4 Antibody.



Western blot analysis of extracts from HeLa cells or recombinant Human H4 using Histone H4 Antibody.

Storage: Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

*Species cross-reactivity is determined by western blot.

**Anti-rabbit secondary antibodies must be used to detect this antibody.

Recommended Antibody Dilutions:

Western Blotting	1:1000
Immunoprecipitation	1:50

For application specific protocols please see the web page for this product at www.cellsignal.com.

Please visit www.cellsignal.com for a complete listing of recommended companion products.

Background References:

- (1) Workman, J.L. and Kingston, R.E. (1998) *Annu. Rev. Biochem.* 67, 545-579.
- (2) Hansen, J.C. et al. (1998) *Biochemistry* 37, 17637-17641.
- (3) Strahl, B.D. and Allis, C.D. (2000) *Nature* 403, 41-45.
- (4) Cheung, P. et al. (2000) *Cell* 103, 263-271.
- (5) Bernstein, B.E. and Schreiber, S.L. (2002) *Chem. Biol.* 9, 1167-1173.
- (6) Jaskelioff, M. and Peterson, C.L. (2003) *Nat. Cell Biol.* 5, 395-399.
- (7) Thorne, A.W. et al. (1990) *Eur. J. Biochem.* 193, 701-713.
- (8) Hendzel, M.J. et al. (1997) *Chromosoma* 106, 348-360.
- (9) Goto, H. et al. (1999) *J. Biol. Chem.* 274, 25543-25549.
- (10) Preuss, U. et al. (2003) *Nucleic Acids Res.* 31, 878-885.
- (11) Dai, J. et al. (2005) *Genes Dev.* 19, 472-488.

IMPORTANT: For western blots, incubate membrane with diluted antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Tween® is a registered trademark of ICI Americas, Inc.