

#2594 Store at -20°C

Acetyl-Histone H4 (Lys8) Antibody



Orders ■ 877-616-CELL (2355)
orders@cellsignal.com
Support ■ 877-678-TECH (8324)
info@cellsignal.com
Web ■ www.cellsignal.com

rev. 02/16/17


For Research Use Only. Not For Use In Diagnostic Procedures.

Applications	Species Cross-Reactivity*	Molecular Wt.	Source
W Endogenous	H, M, R, Mk, (Ce)	11 kDa	Rabbit**

Background: Modulation of chromatin structure plays an important role in the regulation of transcription in eukaryotes. The nucleosome, made up of four core histone proteins (H2A, H2B, H3 and H4), is the primary building block of chromatin (1). The amino-terminal tails of core histones undergo various post-translational modifications, including acetylation, phosphorylation, methylation and ubiquitination (2-5). These modifications occur in response to various stimuli and have a direct effect on the accessibility of chromatin to transcription factors and, therefore, on gene expression (6). In most species, histone H2B is primarily acetylated at lysines 5, 12, 15 and 20 (4,7). Histone H3 is primarily acetylated at Lys9, 14, 18 and 23 (2,3). Acetylation at Lys9 appears to have a dominant role in histone deposition and chromatin assembly in some organisms (2,3). Phosphorylation at Ser10, Ser28 and Thr11 of histone H3 is tightly correlated with chromosome condensation during both mitosis and meiosis (8-10). Phosphorylation of Thr3 of histone H3 is highly conserved among many species and is catalyzed by the kinase haspin. Immunostaining with phospho-specific antibodies in mammalian cells reveals mitotic phosphorylation of H3 Thr3 in prophase and its dephosphorylation during anaphase (11).

Specificity/Sensitivity: Acetyl-Histone H4 (Lys8) Antibody detects endogenous levels of histone H4 only when acetylated at Lys8. The antibody does not cross-react with other acetylated histones.

Source/Purification: Polyclonal antibodies are produced by immunizing animals with a synthetic acetylated peptide corresponding to residues surrounding Lys8 of human histone H4. Antibodies are purified by protein A and peptide affinity chromatography.



Western blot analysis of extracts from various cell lines, untreated or TSA-treated (400 nM TSA for 12 hours), using Acetyl-Histone H4 (Lys8) Antibody.

Entrez-Gene ID #121504
Swiss-Prot Acc. #P62805

Storage: Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

***Species cross-reactivity is determined by western blot.**

****Anti-rabbit secondary antibodies must be used to detect this antibody.**

Recommended Antibody Dilutions:

Western blotting 1:1000

For application specific protocols please see the web page for this product at www.cellsignal.com.

Please visit www.cellsignal.com for a complete listing of recommended companion products.

Background References:

- (1) Workman, J.L. and Kingston, R.E. (1998) *Annu. Rev. Biochem.* 67, 545-579.
- (2) Hansen, J.C. et al. (1998) *Biochemistry* 37, 17637-17641.
- (3) Strahl, B.D. and Allis, C.D. (2000) *Nature* 403, 41-45.
- (4) Cheung, P. et al. (2000) *Cell* 103, 263-271.
- (5) Bernstein, B.E. and Schreiber, S.L. (2002) *Chem. Biol.* 9, 1167-1173.
- (6) Jaskelioff, M. and Peterson, C.L. (2003) *Nat. Cell Biol.* 5, 395-399.
- (7) Thorne, A.W. et al. (1990) *Eur. J. Biochem.* 193, 701-713.
- (8) Hendzel, M.J. et al. (1997) *Chromosoma* 106, 348-360.
- (9) Goto, H. et al. (1999) *J. Biol. Chem.* 274, 25543-25549.
- (10) Preuss, U. et al. (2003) *Nucleic Acids Res.* 31, 878-885.
- (11) Dai, J. et al. (2005) *Genes Dev.* 19, 472-488.

IMPORTANT: For western blots, incubate membrane with diluted antibody in 5% w/v BSA, 1X TBS, 0.1% Tween-20 at 4°C with gentle shaking, overnight.

© 2012 Cell Signaling Technology, Inc. Cell Signaling Technology® is a trademark of Cell Signaling Technology, Inc.

Applications Key: W—Western IP—Immunoprecipitation IHC—Immunohistochemistry ChIP—Chromatin Immunoprecipitation IF—Immunofluorescence F—Flow cytometry E-P—ELISA-Peptide
Species Cross-Reactivity Key: H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken Dm—D. melanogaster X—Xenopus Z—zebrafish B—bovine
 Dg—dog Pg—pig Sc—S. cerevisiae Ce—C. elegans Hr—Horse All—all species expected Species enclosed in parentheses are predicted to react based on 100% homology.