



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

CK2α Antibody

Store at -20C
#2656

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W	Reactivity: H M R Mk	Sensitivity: Endogenous	MW (kDa): 42	Source/Isotype: Rabbit	UniProt ID: #P68400	Entrez-Gene Id: 1457
---------------------------	--------------------------------	-----------------------------------	------------------------	----------------------------------	-------------------------------	--------------------------------

Product Usage Information

Application

Western Blotting

Dilution

1:1000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

This antibody detects endogenous levels of total CK2α 1 protein. This antibody may cross-react with CK2α prime.

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to amino acid residues near the carboxy-terminus of human CK2α1. Antibodies are purified by protein A and peptide affinity chromatography.

Background

CK2 (formerly called Casein Kinase II) is a highly conserved protein kinase with more than 300 substrates regulating cell growth, cell death, and cell survival. CK2 has been implicated in the response to UV irradiation-induced DNA damage, targeting XRCC1 (1) and BRCA1 (2) as well as regulating p53 tumor suppressor protein functions (3). Furthermore, CK2 plays a key role in NF-κB activation (4). UV irradiation stimulates CK2-mediated phosphorylation of several carboxy-terminal residues within IκBα, resulting in IκBα proteasomal degradation and the release and nuclear translocation of active NF-κB. CK2 is also dysregulated in many cancers (5) and neurodegenerative diseases such as Alzheimer's and Parkinson's diseases (6). Structurally, CK2 is a multimeric protein complex consisting of two catalytic subunits (α or α') and two regulatory β subunits (7). CK2 is distributed ubiquitously and is apparently constitutively active (7). While cell cycle-dependent Ser-Pro phosphorylation sites have been identified on CK2α and CK2β, Tyr255 phosphorylation by the Src-related kinase c-Fgr seems to have the greatest effect on CK2α activity (8,9).

Background References

- Morales, J.C. and Carpenter, P.B. (2004) *Sci Aging Knowledge Environ* 2004, pe24.
- O'Brien, K.A. et al. (1999) *Biochem Biophys Res Commun* 260, 658-64.
- Cox, M.L. and Meek, D.W. (2010) *Cell Signal* 22, 564-71.
- Dominguez, I. et al. (2009) *Cell Mol Life Sci* 66, 1850-7.
- Trembley, J.H. et al. *Biofactors* 36, 187-95.
- Perez, D.I. et al. (2010) *Med Res Rev*, Epub ahead of print.
- Bosc, D.G. et al. (1995) *J Biol Chem* 270, 25872-8.
- Donella-Deana, A. et al. (2003) *Biochem J* 372, 841-9.
- Litchfield, D.W. (2003) *Biochem J* 369, 1-15.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting

Cross-Reactivity Key

H: Human **M:** Mouse **R:** Rat **Mk:** Monkey

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's

terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.