

Store at  
-20C  
#26730**HERPUD1 Antibody**

**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

**For Research Use Only. Not for Use in Diagnostic Procedures.**

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W	H M R Mk	Endogenous	54	Rabbit	#Q15011	9709

**Product Usage Information****Application**

Western Blotting

**Dilution**

1:1000

**Storage**

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

**Specificity/Sensitivity**

HERPUD1 Antibody recognizes endogenous levels of total HERPUD1 protein. This antibody also cross-reacts with unidentified proteins of 30 kDa and 48 kDa. This antibody does not cross-react with HERPUD2 protein.

**Source / Purification**

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues near the amino terminus of human HERPUD1 protein. Antibodies are purified by protein A and peptide affinity chromatography.

**Background**

The homocysteine-responsive, ER-resident ubiquitin-like domain member 1 protein (HERPUD1) is an ER membrane protein and putative component of the ER-membrane-associated protein degradation (ERAD) pathway (1). HERPUD1 is a transmembrane protein with both amino- and carboxy-termini exposed to the cytoplasm. The amino-terminal HERPUD1 region contains an ubiquitin-like (UBL) domain (1). Increased expression of HERPUD1 protein following ER stress and its association with ERAD component proteins suggests that HERPUD1 may play an integral role in ERAD pathway function (1). Research studies demonstrate that HERPUD1 associates with components of the ERAD machinery, such as the E3 ubiquitin ligase HRD1, which enhances HRD1-mediated ubiquitination and proteasomal degradation of potentially toxic ERAD substrates (2-4). The HERPUD1 protein may act as a molecular adaptor for the recruitment of ERAD machinery and the amino-terminal UBL domain may important for this function (3,4). The loss of HERPUD1 expression renders cells more susceptible to ER stress and apoptosis (5,6).

**Background References**

1. Kokame, K. et al. (2000) *J Biol Chem* 275, 32846-53.
2. Schulze, A. et al. (2005) *J Mol Biol* 354, 1021-7.
3. Kny, M. et al. (2011) *J Biol Chem* 286, 5151-6.
4. Leitman, J. et al. (2014) *Mol Biol Cell* 25, 1050-60.
5. Hori, O. et al. (2004) *Genes Cells* 9, 457-69.
6. Chan, S.L. et al. (2004) *J Biol Chem* 279, 28733-43.

**Species Reactivity**

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

**Western Blot Buffer**

**IMPORTANT:** For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween@ 20 at 4°C with gentle shaking, overnight.

**Applications Key**

**W:** Western Blotting

**Cross-Reactivity Key**

**H:** Human **M:** Mouse **R:** Rat **Mk:** Monkey

**Trademarks and Patents**

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

**Limited Uses**

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless

separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.