

β-Galactosidase (E2U2I) Rabbit mAb

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W, IF-IC	H M R Hm Mk	Endogenous	65	Rabbit IgG	#P16278	2720
Product Usage Information	Application					Dilution
	Western Blotting					1:1000
	Immunofluorescence (Immunocytochemistry)					1:400
Storage	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.					
Specificity/Sensitivity	β-Galactosidase (E2U2I) Rabbit mAb recognizes endogenous levels of total β-galactosidase protein. The antibody cross-reacts with a 40 kDa protein of unknown identity in some cell extracts.					
Source / Purification	Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Asp39 of human β-galactosidase protein.					
Background	β-galactosidase (also known as β-gal) is an essential hydrolase enzyme that catalyzes the hydrolysis of galactose-containing carbohydrates into monosaccharides. Substrates of β-galactosides include lactose, various glycoproteins, ganglioside GM1, and lactosylceramides. β-galactosidase is used widely in molecular biology; for example, isolation of recombinant bacteria during molecular cloning utilizes α-complementation of the bacterial β-galactosidase gene (<i>lacZ</i>) in the presence of a β-gal substrate to identify recombinant clones (1). In cell biology, Senescence-Associated beta-galactosidase (SA-β-gal), defined as β-gal activity at pH 6.0, is a widely used marker of replicative senescence. While initially thought to derive from a unique isoform of β-galactosidase expressed specifically in senescent cells (2), SA-β-gal activity was subsequently shown to result from overexpression and accumulation of β-galactosidase in endogenous lysosomes, and is not specifically required for replicative senescence (3).					
Background References	<ol style="list-style-type: none"> Messing, J. et al. (1977) <i>Proc Natl Acad Sci U S A</i> 74, 3642-6. Dimri, G.P. et al. (1995) <i>Proc Natl Acad Sci U S A</i> 92, 9363-7. Lee, B.Y. et al. (2006) <i>Aging Cell</i> 5, 187-95. 					

Species Reactivity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).
Western Blot Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.
Applications Key	W: Western Blotting IF-IC: Immunofluorescence (Immunocytochemistry)
Cross-Reactivity Key	H: Human M: Mouse R: Rat Hm: Hamster Mk: Monkey
Trademarks and Patents	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.
Limited Uses	Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect. Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for

diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.