| • | _  | ۰  | ٠ | _  |
|---|----|----|---|----|
|   |    |    |   |    |
|   | ľ  | t  | ī |    |
|   | •  | Ξ  | 1 |    |
|   | F  | ₹  | 1 |    |
|   | ١  | r  | ч |    |
|   | ٠  |    | , |    |
|   |    | ľ  |   |    |
|   |    | Р  | H |    |
|   |    | ١  | J |    |
|   |    | ï  | 7 |    |
|   | ٠  | K  | 4 |    |
|   | ľ  | ٧, | 1 |    |
|   |    | 1  | ï |    |
|   |    |    |   |    |
|   |    |    |   |    |
|   |    |    |   |    |
|   |    |    |   |    |
|   |    | •  | 2 | ١  |
| ı | l  | L  |   | )) |
|   | Z  |    | ı | П  |
| ı | V  | ľ  | ١ | 1  |
|   | ŀ  | ٠  | ١ |    |
|   |    | ١  | i |    |
| ı | ŕ  | 1  | ŧ | П  |
| ı | V. |    | v | ч  |

## **Hip Antibody**



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

## For Research Use Only. Not for Use in Diagnostic Procedures.

| Applications:<br>W           | <b>Reactivity:</b><br>H M R Mk | <b>Sensitivity:</b><br>Endogenous   | MW (kDa):<br>48 | Source/Isotype:<br>Rabbit | UniProt ID:<br>#P50502 | Entrez-Gene Id:<br>6767 |  |  |  |
|------------------------------|--------------------------------|---|-----------------|---------------------------|------------------------|-------------------------|--|--|--|
| Product Usage<br>Information |                                | Application Western Blotting  | 40              | Kassic                    | Dilution<br>1:1000     | 0707                    |  |  |  |
| Storage                      |                                | Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 $\mu$ g/ml BSA and 50% glycerol. Store at – 20°C. Do not aliquot the antibody.  |                 |                           |                        |                         |  |  |  |
| Specificity/Sensitivity      |                                | Hip Antibody detects endogenous levels of total Hip protein.  |                 |                           |                        |                         |  |  |  |
| Source / Purification        |                                | Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to human Hip. Antibodies are purified by protein A and peptide affinity chromatography.   |                 |                           |                        |                         |  |  |  |
| Background                   |                                | Hip (HSP70-interacting protein), also known as ST13 (suppression of tumorigenicity protein 13), is one of several co-chaperones that regulate activities of the HSP70 chaperone family (1,2). The homoligomeric protein Hip cooperates with HSP70 in protein folding by stabilizing the ADP-bound state of HSP70. Hip directly binds to the ATPase domain of HSP70 when it is converted to the ADP-bound state by proteins of the HSP40 family (3). By collaborating with other positive co-factors such as HSP40 and Hop, or competing with negative co-factors such as Bag1, Hip may facilitate the chaperone function of HSP70 in protein folding and repair, and in controlling the activity of regulatory proteins such as steroid receptors and various regulators of proliferation or apoptosis (4-8). |                 |                           |                        |                         |  |  |  |
| Background References        |                                | <ol> <li>Prapapanich, V. et al. (1996) Mol. Endocrino.l 10, 420-431.</li> <li>Gebauer, M. et al. (1997) FEBS Lett. 417, 109-113.</li> <li>Höhfeld, J. et al. (1995) Cell 83, 58998.</li> <li>Frydman, J. and Höhfeld, J. (1997) Trends Biochem. Sci. 22, 87-92.</li> <li>Nollen, E.A. et al. (2001) J. Biol. Chem. 276, 4677-4682.</li> <li>Fan, G.H. et al. (2002) J. Biol. Chem. 277, 6590-6597.</li> <li>Nelson, G.M. et al. (2004) Mol. Endocrinol. 18, 1620-1630.</li> <li>Shi, Z.Z. et al. (2007) J. Zhejiang Univ. Sci. B. 8, 170-176.</li> </ol>  |                 |                           |                        |                         |  |  |  |
|                              |                                |   |                 |                           |                        |                         |  |  |  |

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

**Western Blot Buffer** 

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting

Cross-Reactivity Key

**H:** Human **M:** Mouse **R:** Rat **Mk:** Monkey

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

**Limited Uses** 

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in

any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.