



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Store at -20C
#2771

DYRK1A Antibody

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, IP	Reactivity: H M	Sensitivity: Endogenous	MW (kDa): 90	Source/Isotype: Rabbit	UniProt ID: #Q13627	Entrez-Gene Id: 1859
-------------------------------	---------------------------	-----------------------------------	------------------------	----------------------------------	-------------------------------	--------------------------------

Product Usage Information

Application

Western Blotting
Immunoprecipitation

Dilution

1:1000
1:50

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

DYRK1A Antibody detects endogenous levels of total DYRK1A protein.

Species predicted to react based on 100% sequence homology

Rat

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to human DYRK1A. Antibodies are purified by protein A and peptide affinity chromatography.

Background

The DYRK family includes several dual-specificity tyrosine-phosphorylated and regulated kinases capable of phosphorylating proteins at both Tyr and Ser/Thr residues (1). The DYRK family was identified based on homology to the yeast Yak1 (2) and the *Drosophila* minibrain (mnb) kinases (3). Seven mammalian isoforms have been discovered, including DYRK1A, DYRK1B, DYRK1C, DYRK2, DYRK3, DYRK4, and DYRK4B. Differences in substrate specificity, expression, and subcellular localization are seen across the DYRK family (4,5). All DYRK proteins have a Tyr-X-Tyr motif in the catalytic domain activation loop; phosphorylation of the second Tyr residue (e.g. Tyr312 of DYRK1A) is necessary for kinase activity. DYRKs typically autophosphorylate the Tyr residue within their activation loop, but phosphorylate substrates at Ser and Thr residues (1,6).

DYRK1A phosphorylates serine and threonine residues within a RPX(S/T)P consensus sequence. Substrates include transcription factors such as FoxO1 and cAMP response element-binding proteins such as NFAT (7,8). DYRK1A is ubiquitously expressed in fetal and adult tissues. Transgenic mice with multiple copies of DYRK1A exhibit learning and motor defects suggesting that it is a dosage-sensitive gene (9). The DYRK1A gene localizes to chromosome 21q22.2, a region implicated in Down syndrome, and may contribute to pathological traits observed in chromosome 21 trisomy (10).

Background References

1. Becker, W. and Joost, H.G. (1999) *Prog. Nucleic Acid Res. Mol. Biol.* 62, 1-17.
2. Garrett, S. and Broach, J. (1989) *Genes Dev.* 3, 1336-1348.
3. Tejedor, F. et al. (1995) *Neuron* 14, 287-301.
4. Kentrup, H. et al. (1996) *J. Biol. Chem.* 271, 3488-3495.
5. Becker, W. et al. (1998) *J. Biol. Chem.* 273, 25893-25902.
6. Lochhead, P.A. et al. (2005) *Cell* 121, 925-936.
7. von Groote-Bidlingmaier, F. et al. (2003) *Biochem. Biophys. Res. Commun.* 300, 764-769.
8. Gwack, Y. et al. (2006) *Nature* 441, 646-650.
9. Altafaj, X. et al. (2001) *Hum. Mol. Genet.* 10, 1915-1923.
10. Guimera, J. et al. (1999) *Genomics* 57, 407-418.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween@ 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IP:** Immunoprecipitation

Cross-Reactivity Key

H: Human **M:** Mouse

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.