



**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Store at -20C  
#2785

## Phospho-FADD (Ser191) Antibody (Mouse Specific)

**For Research Use Only. Not for Use in Diagnostic Procedures.**

<b>Applications:</b> W	<b>Reactivity:</b> M	<b>Sensitivity:</b> Endogenous	<b>MW (kDa):</b> 30	<b>Source/Isotype:</b> Rabbit	<b>UniProt ID:</b> #Q61160	<b>Entrez-Gene Id:</b> 14082
---------------------------	-------------------------	-----------------------------------	------------------------	----------------------------------	-------------------------------	---------------------------------

### Product Usage Information

#### Application

Western Blotting

#### Dilution

1:1000

### Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

### Specificity/Sensitivity

Phospho-FADD (Ser191) Antibody (Mouse Specific) detects endogenous levels of mouse FADD protein only when phosphorylated at serine 191.

### Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Ser191 of mouse FADD. Antibodies are purified by protein A and peptide affinity chromatography.

### Background

Fas-associated death domain (FADD or Mort 1) functions as an important adaptor in coupling death signaling from membrane receptors, such as the Fas ligand and TNF family (DR3, DR4 and DR5), to caspase-8 (1,2). FADD has a carboxy-terminal death domain, which interacts with the cytoplasmic tail of the membrane receptor, and an amino-terminal death effector domain, which interacts with caspase-8. Clustering of the receptors upon stimulation brings about FADD and caspase-8 oligomerization, activating the caspase signaling pathway. Human FADD is phosphorylated mainly at Ser194, while mouse FADD is phosphorylated at Ser191. In both cases, the phosphorylation is cell cycle-dependent (3) and may be related to its regulatory role in embryonic development and cell cycle progression (4,5).

### Background References

- Ashkenazi, A. and Dixit, V.M. (1998) *Science* 281, 1305-1308.
- Kuang, A. A. et al. (2000) *J. Biol. Chem.* 275, 25065-25068.
- Scaffidi, C. et al. (2000) *J. Immunol.* 164, 1236-1242.
- Newton, K. et al. (2000) *EMBO J.* 19, 931-941.
- Zhang, J. et al. (2001) *J. Biol. Chem.* 276, 29815-29818.

### Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

### Western Blot Buffer

**IMPORTANT:** For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween@ 20 at 4°C with gentle shaking, overnight.

### Applications Key

**W:** Western Blotting

### Cross-Reactivity Key

**M:** Mouse

### Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

### Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for

diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.