

Store at
-20°C

Adiponectin (C45B10) Rabbit mAb

www.cellsignal.com

#2789

Support: 877-678-TECH (8324)
www.cellsignal.com/support**Orders:** 877-616-CELL (2355)
orders@cellsignal.com**Entrez-Gene ID** #9370
UniProt ID #Q15848

rev. 10/03/17

For Research Use Only. Not For Use In Diagnostic Procedures.

Applications W Endogenous	Species Cross-Reactivity* H, M, R	Molecular Wt. 27 kDa	Isotype Rabbit IgG**
---------------------------------	--------------------------------------	-------------------------	-------------------------


Background: Adiponectin, also termed AdipoQ, Acrp30, apM1 and GBP28, is an adipokine expressed exclusively in brown and white adipocytes (1). It is secreted into the blood and exists in three major forms: a low molecular weight trimer, a medium molecular weight hexamer and a high molecular weight multimer (1). Adiponectin levels are decreased in obese and insulin-resistant mice and humans (2), suggesting that this adipokine is critical to maintain insulin sensitivity. Adiponectin stimulates the phosphorylation of AMPK α at Thr172 and activates AMPK in skeletal muscle (3). It also stimulates glucose uptake in myocytes (3). The block of AMPK activation by a dominant-negative AMPK α 2 isoform inhibits the effect of adiponectin on glucose uptake, indicating that adiponectin stimulates glucose uptake and increases insulin sensitivity through its action on AMPK (3). Adiponectin mutants that are not able to form oligomers larger than trimers have no effect on the AMPK pathway (4). Mutations that render adiponectin unable to form high molecular weight multimers are associated with human diabetes (4), indicating the importance of multimerization for adiponectin activity.

Specificity/Sensitivity: Adiponectin (C45B10) Rabbit mAb detects endogenous levels of total adiponectin protein monomer. It will not detect higher molecular weight forms of adiponectin.

Source/Purification: Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to human adiponectin.

Background References:

- (1) Kadowaki, T. et al. (2006) *J. Clin. Invest.* 116, 1784-1792.
- (2) Hu, E. et al. (1996) *J. Biol. Chem.* 271, 10697-10703.
- (3) Yamauchi, T. et al. (2002) *Nat. Med.* 8, 1288-1295.
- (4) Waki, H. et al. (2003) *J. Biol. Chem.* 278, 40352-40363.



Western blot analysis of extracts from NIH/3T3 and 3T3-L1 cells using Adiponectin (C45B10) Rabbit mAb.

Storage: Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C . Do not aliquot the antibody.

*Species cross-reactivity is determined by western blot.

**Anti-rabbit secondary antibodies must be used to detect this antibody.

Recommended Antibody Dilutions:

Western blotting 1:1000

For product specific protocols and a complete listing of recommended companion products please see the product web page at www.cellsignal.com

U. S. Patent No. 5,675,063
Tween[®] is a registered trademark of ICI Americas, Inc.

Thank you for your recent purchase. If you would like to provide a review visit www.cellsignal.com/comments.

IMPORTANT: For western blots, incubate membrane with diluted antibody in 5% w/v BSA, 1X TBS, 0.1% Tween[®]20 at 4°C with gentle shaking, overnight.

© 2014 Cell Signaling Technology, Inc.
Cell Signaling Technology[®] is a trademark of Cell Signaling Technology, Inc.



Applications: W—Western IP—Immunoprecipitation IHC—Immunohistochemistry ChIP—Chromatin Immunoprecipitation IF—Immunofluorescence F—Flow cytometry E-P—ELISA-Peptide Species Cross-Reactivity: H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken Dm—D. melanogaster X—Xenopus Z—zebrafish B—bovine Dg—dog Pg—pig Sc—S. cerevisiae Ce—C. elegans Hr—Horse All—all species expected Species enclosed in parentheses are predicted to react based on 100% homology.