



Orders:	877-616-CELL (2355) orders@cellsignal.com
Support:	877-678-TECH (8324)
Web:	info@cellsignal.com cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, W-S, IP	<b>Reactivity:</b> H M R Hm Mk	<b>Sensitivity:</b> Endogenous	<b>MW (kDa):</b> 34	<b>Source/Isotype:</b> Rabbit IgG	<b>UniProt ID:</b> #P06493	Entrez-Gene Id: 983		
Product Usage Information	e	<b>Application</b> Western Blotting Simple Western™ Immunoprecipitation			<b>Dilution</b> 1:1000 1:10 - 1:50 1:50			
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.						
Specificity/Sensitivity		Cdc2 (E1Z6R) Rabbit mAb recognizes endogenous levels of total cdc2 protein.						
Source / Purification		Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Asp289 of human cdc2 protein.						
Background		The entry of eukaryotic cells into mitosis is regulated by cdc2 kinase activation, a process controlled at several steps including cyclin binding and phosphorylation of cdc2 at Thr161 (1). However, the critical regulatory step in activating cdc2 during progression into mitosis appears to be dephosphorylation of cdc2 at Thr14 and Tyr15 (2). Phosphorylation at Thr14 and Tyr15, resulting in inhibition of cdc2, can be carried out by Wee1 and Myt1 protein kinases (3,4). The cdc25 phosphatase may be responsible for removal of phosphates at Thr14 and Tyr15 and subsequent activation of cdc2 (1,5).						
Background R	eferences	1. Atherton-Fessler, S. 2. Norbury, C. et al. (19 3. McGowan, C.H. and 4. Wells, N.J. et al. (199 5. Hunter, T. (1995) <i>Ce</i> .	991) <i>EMBO J</i> 10, 332 Russell, P. (1993) <i>E</i> 99) <i>J Cell Sci</i> 112 ( Pt	21-9. <i>MBO J</i> 12, 75-85.				
Species React	ivity	Species reactivity is de	etermined by testing	g in at least one approve	ed application (e.g.,	western blot).		
Western Blot	Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.				n 5% w/v BSA, 1X		
Applications <b>F</b>	(ey	W: Western Blotting W-S: Simple Western™ IP: Immunoprecipitation						
Cross-Reactiv	ity Key	H: Human M: Mouse R: Rat Hm: Hamster Mk: Monkey						
Trademarks a	nd Patents	Patents Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.						
		XP is a registered trademark of Cell Signaling Technology, Inc.						
		All other trademarks a more information.	are the property of t	heir respective owners.	Visit cellsignal.com	/trademarks for		
Limited Uses		the following terms ap terms and conditions	oply to Products pro that are in addition	a writing signed by a leg ovided by CST, its affiliat to, or different from, th y authorized representa	es or its distributors ose contained here	s. Any Customer's in, unless		
		approved, cleared, or purpose. Customer sh any manner that confl	licensed by the FDA all not use any Prod licts with its labeling	se Only or a similar labe or other regulatory for duct for any diagnostic o g statement. Products so research and developme	eign or domestic er or therapeutic purp old or licensed by C	itity, for any ose, or otherwise in ST are provided for		

diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.