

#2862 Store at -20°C

c-Abl Antibody



Orders ■ 877-616-CELL (2355)
orders@cellsignal.com
Support ■ 877-678-TECH (8324)
info@cellsignal.com
Web ■ www.cellsignal.com

rev. 10/13/16

For Research Use Only. Not For Use In Diagnostic Procedures.

Applications	Species Cross-Reactivity*	Molecular Wt.	Source
W, IP Endogenous	H, M, R	135 (c-Abl) 210 (Bcr-Abl)	Rabbit**


Background: The c-Abl proto-oncogene encodes a nonreceptor protein tyrosine kinase that is ubiquitously expressed and highly conserved in metazoan evolution. c-Abl protein is distributed in both the nucleus and the cytoplasm of cells. It is implicated in regulating cell proliferation, differentiation, apoptosis, cell adhesion and stress responses (1-3). c-Abl kinase activity is increased *in vivo* by diverse physiological stimuli including integrin activation, PDGF stimulation and binding to c-Jun, Nck and RFX1 (2,4). The *in vivo* mechanism of regulation of c-Abl kinase activity is not completely understood. Tyr245 is located in the linker region between the SH2 and catalytic domains. This positioning is conserved among Abl family members. Phosphorylation of Tyr245 is involved in the activation of c-Abl kinase (5). In addition, phosphorylation of Tyr412, which is located in the kinase activation loop of c-Abl, is required for kinase activity (6).

Specificity/Sensitivity: c-Abl Antibody detects endogenous levels of Abl proteins.

Source/Purification: Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to the sequence close to the carboxy-terminus of human c-Abl. Antibodies are purified by protein A and peptide affinity chromatography.

Background References:

- (1) Wang, J.Y. et al. (2000) *Oncogene* 19, 5643–5650.
- (2) Van Etten, R.A. et al. (1999) *Trends Cell. Biol.* 9, 179–182.
- (3) Danial, N.N. et al. (2000) *Oncogene* 19, 2523–2531.
- (4) Shaul, Y. et al. (2000) *Cell Death Differ.* 7, 10–16.
- (5) Brasher, B.B. et al. (2000) *J. Biol. Chem.* 275, 35631–35637.
- (6) Pluk, H. et al. (2002) *Cell* 108, 247–259.



Western blot analysis of extracts from various cell lines, using c-Abl Antibody.

Entrez-Gene ID #25
Swiss-Prot Acc. #P00519-2

Storage: Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

***Species cross-reactivity is determined by western blot.**
****Anti-rabbit secondary antibodies must be used to detect this antibody.**

Recommended Antibody Dilutions:

Western blotting 1:1000
Immunoprecipitation 1:100

For application specific protocols please see the web page for this product at www.cellsignal.com.

Please visit www.cellsignal.com for a complete listing of recommended companion products.

IMPORTANT: For western blots, incubate membrane with diluted antibody in 5% w/v BSA, 1X TBS, 0.1% Tween-20 at 4°C with gentle shaking, overnight.

DRAQ5® is a registered trademark of Biostatus Limited.

© 2011 Cell Signaling Technology, Inc. Cell Signaling Technology® is a trademark of Cell Signaling Technology, Inc.

Applications Key: W—Western IP—Immunoprecipitation IHC—Immunohistochemistry ChIP—Chromatin Immunoprecipitation IF—Immunofluorescence F—Flow cytometry E-P—ELISA-Peptide
Species Cross-Reactivity Key: H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken Dm—D. melanogaster X—Xenopus Z—zebrafish B—bovine
 Dg—dog Pg—pig Sc—S. cerevisiae Ce—C. elegans Hr—Horse All—all species expected Species enclosed in parentheses are predicted to react based on 100% homology.