



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

#2877 Store at -20C

FE65 Antibody

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W	Reactivity: M R	Sensitivity: Endogenous	MW (kDa): 100	Source/Isotype: Rabbit	UniProt ID: #O00213	Entrez-Gene Id: 322
---------------------------	---------------------------	-----------------------------------	-------------------------	----------------------------------	-------------------------------	-------------------------------

Product Usage Information

Application

Western Blotting

Dilution

1:1000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

FE65 Antibody detects endogenous levels of FE65. It does not cross-react with FE65L1 or FE65L2.

Species predicted to react based on 100% sequence homology

Human

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues of human FE65. Antibodies are purified by protein A and peptide affinity chromatography.

Background

FE65, FE65L1, and FE65L2 are members of the FE65 protein family. FE65 is an adaptor protein with protein-protein interaction domains, including a WW domain followed by two phosphotyrosine interaction domains (PID1 and PID2) (1). Amyloid-β (Aβ) precursor protein (APP) binds to PID2 and undergoes sequential cleavage. First, α-/β secretases cleave and release the ectodomain into the extracellular environment. Subsequent processing by the γ-secretase complex results in the APP intracellular domain (AICD) and the Aβ peptides. The latter Aβ fragments form the main components of amyloid plaques in patients with Alzheimer's disease (2). FE65 family members can regulate APP processing, resulting in elevated levels of Aβ (3). Double knockout mice of FE65 and FE65L1 display a phenotype that occurs in animals lacking APP family members, supporting a functional interaction between FE65 and APP (4).

Background References

1. Russo, T. et al. (1998) *FEBS Lett* 434, 1-7.
2. Selkoe, D.J. (1996) *J Biol Chem* 271, 18295-8.
3. King, G.D. and Scott Turner, R. (2004) *Exp Neurol* 185, 208-19.
4. Guénette, S. et al. (2006) *EMBO J* 25, 420-31.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting

Cross-Reactivity Key

M: Mouse **R:** Rat

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any

purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.