

Cyclin H Antibody



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W, IP	H M R	Endogenous	36	Rabbit	#P51946	902

Product Usage Information

Application

Western Blotting
Immunoprecipitation

Dilution

1:1000
1:100

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

Cyclin H Antibody detects endogenous levels of cyclin H. It does not cross-react with other family members at physiological levels.

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues near the carboxy-terminus of cyclin H. Antibodies are purified by protein A and peptide affinity chromatography.

Background

Cyclin H belongs to a conserved cyclin family that plays a critical role in the regulation of cell cycle dependent kinases (CDKs) necessary for cell cycle progression (1,2). In general, the activity of CDKs requires the binding of appropriate cyclins as well as phosphorylation driven by Cdk-activating kinase (CAK). Cyclin H is part of the CAK complex that includes the kinase CDK7, and an assembly factor p36/Mat1, which enhances binding between cyclin H and CDK7 and increases activity (3,4). CAK regulates progression through the cell cycle by activating cdc2, CDK2, and CDK4 kinases through phosphorylation of a critical threonine residue in the T-loop of the CDK-cyclin complexes (5,6). The CAK complex can exist either in its free form or in association with transcription factor IIH (TFIIH) which can affect its substrate specificity (7,8,9). When bound to TFIIH, CAK preferentially phosphorylates the carboxy-terminal domain of RNA polymerase II (9), providing a link between cell cycle control, transcriptional regulation, and DNA repair.

Background References

1. Fisher, R.P. and Morgan, D.O. (1994) *Cell* 78, 713-24.
2. Mäkelä, T.P. et al. (1994) *Nature* 371, 254-7.
3. Yee, A. et al. (1995) *Cancer Res* 55, 6058-62.
4. Devault, A. et al. (1995) *EMBO J* 14, 5027-36.
5. Solomon, M.J. (1994) *Trends Biochem Sci* 19, 496-500.
6. Morgan, D.O. (1995) *Nature* 374, 131-4.
7. Shiekhhattar, R. et al. (1995) *Nature* 374, 283-7.
8. Serizawa, H. et al. (1995) *Nature* 374, 280-2.
9. Rossignol, M. et al. (1997) *EMBO J* 16, 1628-37.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IP:** Immunoprecipitation

Cross-Reactivity Key

H: Human **M:** Mouse **R:** Rat

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's

terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.