Erlin-2 Antibody



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, IP	Reactivity: H M	Sensitivity: Endogenous	MW (kDa): 43	Source/Isotype: Rabbit	UniProt ID: #O94905	Entrez-Gene Id: 11160
Product Usage Information		Application Western Blotting Immunoprecipitation			Dilution 1:1000 1:50	
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA and 50% glycerol. Store at – 20°C. Do not aliquot the antibody.				
Specificity/Sensitivity		Erlin-2 Antibody detects endogenous levels of human erlin-2 protein.				
Source / Purification		Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Gln315 of mouse erlin-2. Antibodies are purified by peptide affinity chromatography.				
Background		Erlin-1 and erlin-2 (SPFH1 and SPFH2) are SPFH domain-containing proteins that belong to the prohibitin family (1,2). The N-termal domain of erlin proteins contains ER-targeting sequences responsible for their translocation to the endoplasmic reticulum (ER) (3). In the ER, erlin-1 and erlin-2 specifically associate with the detergent resistant lipid raft microdomain of the membrane (3). Erlin-1 may be involved in dentritic cell activation (4) and erlin-2 has been shown to regulate the ER-associated degradation (ERAD) pathway by interacting with endogenous substrates and resulting in their polyubiquitination and degradation (5).				
Background References		1. Browman, D.T. et al. (2007) <i>Trends Cell Biol</i> 17, 394-402. 2. Morrow, I.C. and Parton, R.G. (2005) <i>Traffic</i> 6, 725-40. 3. Browman, D.T. et al. (2006) <i>J Cell Sci</i> 119, 3149-60. 4. Li, N. et al. (2000) <i>Biochem Biophys Res Commun</i> 279, 487-93. 5. Pearce, M.M. et al. (2007) <i>J Biol Chem</i> 282, 20104-15.				
Species Reactiv	rity	Species reactivity is de	termined by testing	g in at least one approve	ed application (e.g.,	western blot).
Western Blot Buffer		IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.				
Applications Key		W: Western Blotting IP: Immunoprecipitation				
Cross-Reactivity Key		H: Human M: Mouse				
Trademarks and Patents		Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.				
		All other trademarks a more information.	re the property of t	their respective owners.	Visit cellsignal.com	/trademarks for
Limited Uses		Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.				
		Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any				

purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for

diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.