s 686

Limited Uses

force or effect.

CCL5/RANTES Antibody



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

| Applications: W, IP | Reactivity: M | Sensitivity: Transfected Only | MW (kDa): 10 | Source/Isotype: Rabbit | UniProt ID: #P13501 | Entrez-Gene Id: 6352 |
|--|-------------------------|---|--|--|-----------------------------|-------------------------|
| Product Usage Information | | Application Western Blotting Immunoprecipitation | | | Dilution 1:1000 1:25 | |
| Storage | | Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA and 50% glycerol. Store at – 20°C. Do not aliquot the antibody. | | | | |
| Specificity/Sensitivity | | CCL5/RANTES Antibody detects transfected levels of CCL5/RANTES. | | | | |
| Species predicted to react based on 100% sequence homology | | Rat | | | | |
| Source / Purification | | Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues at the carboxyl terminus of rodent CCL5/RANTES. | | | | |
| Background | | RANTES/CCL5 (regulated upon activation, T cell expressed and secreted) is a member of the C-C or β family of chemokines that induce inflammation and are associated with a number of inflammatory disorders (1,2). RANTES is produced and secreted mainly by CD8 ⁺ T cells, macrophages, and platelets, as well as epithelial cells, fibroblasts, and some solid tumors (2-7). RANTES acts as a chemoattractant and has other regulatory functions on a number of cell types, including monocytes, memory T cells, NK cells, eosinophils, basophils, dendritic cells, and mast cells (3, 7-9). Signaling by RANTES is mediated by several G protein-coupled receptors (GPCRs), including CCR1, CCR3, CCR4, and CCR5. | | | | |
| Background References | | 1. Schall, T.J. (1991) <i>Cyt</i> 2. Schall, T.J. et al. (198 3. Kameyoshi, Y. et al. (4. Rathanaswami, P. et 5. Mattei, S. et al. (1994 6. Stellato, C. et al. (1997 7. Schall, T.J. et al. (1999 8. Kuna, P. et al. (1992) 9. Mattoli, S. et al. (199 | 8) <i>J Immunol</i> 141, (1992) <i>J Exp Med</i> 1 al. (1993) <i>J Biol Ch</i> 4) <i>Int J Cancer</i> 56, 8 95) <i>J Immunol</i> 155, 0) <i>Nature</i> 347, 669 <i>J Immunol</i> 149, 63 | 76, 587-92. em 268, 5834-9. 353-7. 410-8. -71. | 5-21. | |
| Species Reacti | vity | Species reactivity is de | termined by testin | g in at least one approve | ed application (e.g., | western blot). |
| Western Blot Buffer | | IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight. | | | | |
| Applications Key | | W: Western Blotting IP: Immunoprecipitation | | | | |
| Cross-Reactivity Key | | M: Mouse | | | | |
| Trademarks and Patents | | Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. | | | | |
| | | All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information. | | | | |

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.