

NTF2 (5A3) Mouse mAb

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W, IF-IC	H M R Mk	Endogenous	14	Mouse IgG2a	#P61970	10204

Product Usage Information**Application**

Western Blotting
Immunofluorescence (Immunocytochemistry)

Dilution

1:1000
1:50

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

NTF2 (5A3) Mouse mAb detects endogenous levels of total NTF2 protein.

Source / Purification

Monoclonal antibody is produced by immunizing animals with full-length recombinant human NTF2.

Background

The small GTPase Ran resides on both the cytosolic and nucleosolic sides of the nuclear pore complex (NPC) and regulates the import and export of various proteins to and from the nucleus. Like other small GTPases, Ran exists in either a GTP-bound or GDP-bound state. RanGTP that resides in the nucleus and promotes nuclear export, while cytosolic RanGDP promotes import. The gradient of RanGTP across the nuclear membrane allows for appropriate movement of cargo proteins across the NPC as well as maintenance of the mitotic spindle (1-3).

Nuclear transport factor 2 (NTF2) regulates the subcellular distribution and function of Ran (4-5). The NTF2 homodimer facilitates the diffusion of RanGDP through NPCs via transient interactions with phenylalanine-glycine (FG) repeat domains on NPC proteins. NTF2 stabilizes the GDP-bound form of Ran until it is induced to dissociate by a nuclear factor in an ATP-dependent manner, thus allowing the guanine nucleotide exchange factor (GEF) RCC1 to mediate exchange of GDP for GTP on Ran (6-7).

Background References

- Mattaj, I.W. and Englmeier, L. (1998) *Annu Rev Biochem* 67, 265-306.
- Kalab, P. et al. (2002) *Science* 295, 2452-6.
- Becksei, A. and Mattaj, I.W. (2003) *Proc Natl Acad Sci USA* 100, 1717-22.
- Ribbeck, K. et al. (1998) *EMBO J* 17, 6587-98.
- Steggerda, S.M. et al. (2000) *Mol Biol Cell* 11, 703-19.
- Stewart, M. (2000) *Cell Struct Funct* 25, 217-25.
- Yamada, M. et al. (2004) *J Biol Chem* 279, 36228-34.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween@ 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IF-IC:** Immunofluorescence (Immunocytochemistry)

Cross-Reactivity Key

H: Human **M:** Mouse **R:** Rat **Mk:** Monkey

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for

Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.

Orders: 877-616-CELL (2355) • orders@cellsignal.com • Support: 877-678-TECH (8324) • info@cellsignal.com • Web: cellsignal.com
For Research Use Only. Not for Use in Diagnostic Procedures.