3120 Store at -20C

VASP (A290) Antibody



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W	Reactivity: H Mk	Sensitivity: Endogenous	MW (kDa): 46, 50	Source/Isotype: Rabbit	UniProt ID: #P50552	Entrez-Gene Id: 7408
Product Usage Information		Application Western Blotting			Dilution 1:1000	
Storage		Supplied in 10 mM so 20°C. Do not aliquot t), 150 mM NaCl, 100 μg.	/ml BSA and 50% gl	ycerol. Store at –
Specificity/Sensitivity		VASP (A290) Antibody detects endogenous levels of total VASP protein.				
Source / Purification		Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding the carboxy-terminal sequence of human VASP. Antibodies are purified by peptide affinity chromatography.				
Background		Vasodilator-stimulated phosphoprotein (VASP) was originally characterized as a substrate of both cGMP- and cAMP-dependent kinases (PKG and PKA, or cGPK and cAPK, respectively) (1). It is now believed that VASP belongs to the Ena/VASP family of adaptor proteins linking the cytoskeletal system to the signal transduction pathways and that it functions in cytoskeletal organization, fibroblast migration, platelet activation, and axon guidance (2,3). Three phosphorylation sites, Ser157, Ser239, and Thr278, have been identified. Ser239 is the major PKG phosphorylation site, while Ser157 is the major PKA phosphorylation site (4). Evidence suggests that VASP phosphorylation reduces its association with actin and has a negative effect on actin polymerization (5). Phosphorylation at Ser239 of VASP is a useful marker for monitoring PKG activation and signaling (6,7).				
Background References		 Butt, E. et al. (1994) J Biol Chem 269, 14509-17. Ball, L.J. et al. (2000) EMBO J 19, 4903-14. Machesky, L.M. (2000) Cell 101, 685-8. Smolenski, A. et al. (1998) J Biol Chem 273, 20029-35. Harbeck, B. et al. (2000) J Biol Chem 275, 30817-25. Oelze, M. et al. (2000) Circ Res 87, 999-1005. Lawrence, D.W. and Pryzwansky, K.B. (2001) J Immunol 166, 5550-6. 				
_		3. Machesky, L.M. (2000) <i>Cell</i> 101, 685-8. 4. Smolenski, A. et al. (1998) <i>J Biol Chem</i> 273, 20029-35. 5. Harbeck, B. et al. (2000) <i>J Biol Chem</i> 275, 30817-25. 6. Oelze, M. et al. (2000) <i>Circ Res</i> 87, 999-1005.				

Species Reactivity Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X

TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key W: Western Blotting

Cross-Reactivity Key H: Human Mk: Monkey

Trademarks and Patents Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for

more information.

Limited UsesExcept as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to or different from those contained berein, unless

terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no

force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in

any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.