

#3279
Store at -20°C**Shootin1 Antibody**
Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, IP	Reactivity: H M R	Sensitivity: Endogenous	MW (kDa): 60	Source/Isotype: Rabbit	UniProt ID: #A0MZ66	Entrez-Gene Id: 57698
Product Usage Information		Application Western Blotting Immunoprecipitation		Dilution 1:1000 1:100		
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.				
Specificity/Sensitivity		Shootin1 Antibody detects endogenous levels of total shootin1 protein.				
Source / Purification		Polyclonal antibodies are produced by immunizing animals with a synthetic peptide surrounding Lys450 of human shootin1. Antibodies are purified by peptide affinity chromatography.				
Background		The differentiation process of neurons can be divided into five stages, each stage characterized by morphological changes observed in the developing cells. In stage 1, the cells extend lamellipodia and in stage 2 their lamellipodia develop into immature neurites. In stage 3 one neurite elongates rapidly to form an axon and in stage 4 the remaining immature neuritis elongate to form dendrites. In stage 5 synaptic contacts are formed and a neuronal network is established (1). Shootin1 is involved in generating internal asymmetric signals required for neuronal during stages 2 and 3. The extension of an axon requires considerable reorganization of the cytoskeleton mediated by PI3K/Akt and PI3K/Cdc42 signaling (1). Shootin1 is involved in regulating the subcellular localization of PI3 kinase. Furthermore, shootin1 is upregulated during polarization and accumulates asymmetrically in a single neurite that consequently elongates rapidly to form an axon (2).				
Background References		1. Yoshimura, T. et al. (2006) <i>J Neurosci</i> 26, 10626-30. 2. Toriyama, M. et al. (2006) <i>J Cell Biol</i> 175, 147-57.				

Species Reactivity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).
Western Blot Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.
Applications Key	W: Western Blotting IP: Immunoprecipitation
Cross-Reactivity Key	H: Human M: Mouse R: Rat
Trademarks and Patents	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.
Limited Uses	<p>Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.</p> <p>Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any</p>

commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.

Orders: 877-616-CELL (2355) • orders@cellsignal.com • Support: 877-678-TECH (8324) • info@cellsignal.com • Web: cellsignal.com
For Research Use Only. Not for Use in Diagnostic Procedures.