Revision 1

Store at

32961

នុ Inflammasome Antibody Sampler Kit



| Orders: | 877-616-CELL (2355) orders@cellsignal.com |
|----------|--|
| Support: | 877-678-TECH (8324) |
| Web: | info@cellsignal.com cellsignal.com |

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

| Product Includes | Product # | Quantity | Mol. Wt | Isotype/Source |
|---|-----------|----------|----------------|----------------|
| NLRP3 (D4D8T) Rabbit mAb | 15101 | 20 µl | 110 kDa | Rabbit IgG |
| AIM2 (D5X7K) Rabbit mAb | 12948 | 20 µl | 40 kDa | Rabbit IgG |
| NLRC4 (D5Y8E) Rabbit mAb | 12421 | 20 µl | 110 kDa | Rabbit IgG |
| Caspase-1 (D7F10) Rabbit mAb | 3866 | 20 µl | 48, 20 kDa | Rabbit IgG |
| NALP1 Antibody | 4990 | 20 µl | 165, 70 kDa | Rabbit |
| ASC/TMS1 (E1E3I) Rabbit mAb | 13833 | 20 µl | 22, 19, 15 kDa | Rabbit IgG |
| Cleaved Caspase-1 (Asp297) (D57A2) Rabbit mAb | 4199 | 20 µl | 20, 22 kDa | Rabbit IgG |
| Cleaved-IL-1β (Asp116) (D3A3Z) Rabbit mAb | 83186 | 20 µl | 17 kDa | Rabbit IgG |
| IL-1β (D3U3E) Rabbit mAb | 12703 | 20 µl | 17, 31 kDa | Rabbit IgG |
| Anti-rabbit IgG, HRP-linked Antibody | 7074 | 100 µl | | Goat |
| | | | | |

Please visit cellsignal.com for individual component applications, species cross-reactivity, dilutions, protocols, and additional product information.

| Description | The Inflammasome Antibody Sampler Kit provides an economical means of detecting multiple inflammasome components. The kit contains enough primary antibodies to perform at least two western blot experiments. |
|-----------------------|---|
| Storage | Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody. |
| Background | The innate immune system works as the first line of defense in protection from pathogenic microbes and host-derived signals of cellular distress. One way in which these "danger" signals trigger inflammation is through activation of inflammasomes, which are multiprotein complexes that assemble in the cytosol after exposure to pathogen-associated molecular patterns (PAMPs) or danger- associated molecular patterns (DAMPs) and result in the activation of caspase-1 and subsequent cleavage of proinflammatory cytokines IL-1 β and IL-18 (Reviewed in 1-6). Inflammasome complexes typically consist of a cytosolic pattern recognition receptor (PRR; a nucleotide-binding domain and leucine-rich-repeat [NLR] or AIM2-like receptor [ALR] family member), an adaptor protein (ASC/TMS1), and pro-caspase-1. A number of distinct inflammasome complexes have been identified, each with a unique PRR and activation triggers. The best characterized is the NLRP3 complex, which contains NLRP3, ASC/TMS1, and pro-caspase-1. The NLRP3 inflammasome is activated in a two-step process. First, NF-kB signaling is induced through PAMP- or DAMP-mediated activation of TLR4 or TNFR, resulting in increased expression of NLRP3, pro-IL-1 β , and pro-IL-18 (priming step, signal 1). Next, indirect activation of NLRP3 occurs by a multitude of signals (whole pathogens, PAMPs/DAMPs, potassium efflux, lysosomal-damaging environmental factors [uric acid, silica, alum] and endogenous factors [amyloid- β , cholesterol crystals], and mitochondrial damage), leading to complex assembly and activation of caspase-1 (signal 2). The complex inflammasome structure is built via domain interactions among the protein components. Other inflammasomes are activated by more direct means: double- stranded DNA activates the AIM2 complex, anthrax toxin activates NLRP1, and bacterial flagellin activates NLRC4. Activated caspase-1 induces secretion of proinflammatory cytokines IL-1 β and -18, but also regulates metabolic enzyme expression, phagosome maturation, vasodilation, and py |
| Background References | Broz, P. and Dixit, V.M. (2016) Nat Rev Immunol 16, 407-20. Guo, H. et al. (2015) Nat Med 21, 677-87. Jo, E.K. et al. (2016) Cell Mol Immunol 13, 148-59. Rathinam, V.A. and Fitzgerald, K.A. (2016) Cell 165, 792-800. Shao, B.Z. et al. (2015) Front Pharmacol 6, 262. Schroder, K. and Tschopp, J. (2010) Cell 140, 821-32. |

| Trademarks and Patents | Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. | | |
|------------------------|--|--|--|
| | All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information. | | |
| Limited Uses | Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect. | | |
| | Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products. | | |