

**Dab1 Antibody**

**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

**For Research Use Only. Not for Use in Diagnostic Procedures.**

<b>Applications:</b> W	<b>Reactivity:</b> H	<b>Sensitivity:</b> Transfected Only	<b>MW (kDa):</b> 80, 110 GFP-Dab1 fusion.	<b>Source/Isotype:</b> Rabbit	<b>UniProt ID:</b> #O75553	<b>Entrez-Gene Id:</b> 1600
---------------------------	-------------------------	---	--	----------------------------------	-------------------------------	--------------------------------

**Product Usage Information****Application**

Western Blotting

**Dilution**

1:1000

**Storage**

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

**Specificity/Sensitivity**

Dab1 Antibody detects transfected levels of Dab1 protein. This antibody does not cross-react with other proteins.

**Species predicted to react based on 100% sequence homology**

Mouse, Rat

**Source / Purification**

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues close to the amino terminus of human Dab1. Antibodies are purified by protein A and peptide affinity chromatography.

**Background**

The Reelin signaling pathway plays a critical role in neuronal development. Reelin is a secreted glycoprotein that binds to the lipoprotein receptors VLDLR and ApoER2 or alpha3beta1 integrin on the surface of neurons (1,2). Activation of these receptors induces tyrosine phosphorylation of Disabled 1 (Dab1), an intracellular adaptor. It is generally believed that tyrosine phosphorylation of Dab1 by Src family tyrosine kinases is the most critical downstream event in Reelin signaling. The phosphotyrosine-binding (PTB) domain within its amino terminus enables Dab1 to recognize and bind to a conserved sequence motif within the cytoplasmic tail of the receptors. In addition, the PTB contains a Pleckstrin Homology-like subdomain that binds to phosphoinositides. The phosphoinositide-binding region within the Dab1 PTB domain is required for membrane localization and basal tyrosine phosphorylation of Dab1 independent of VLDLR and ApoER2 (3). It has been demonstrated that Src, CrkII, CrkL and Dock1 associate with tyrosine-phosphorylated Dab. The CrkII-Dab1 interaction requires tyrosine phosphorylation of Dab1 at residues 220 or 232 (4).

**Background References**

1. Huang, Y. et al. (2005) *Biochem. Biophys. Res. Commun.* 331, 1460-1468.
2. Luque, J.M. (2004) *Brain Res. Dev. Brain Res.* 152, 269-271.
3. Morimura, T. et al. (2005) *J. Biol. Chem.* 280, 16901-16908.
4. Chen, K. et al. (2004) *J. Cell. Sci.* 117, 4527-4536.

**Species Reactivity**

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

**Western Blot Buffer**

**IMPORTANT:** For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

**Applications Key**

**W:** Western Blotting

**Cross-Reactivity Key**

**H:** Human

**Trademarks and Patents**

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

**Limited Uses**

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's

terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.