

ALK (C26G7) Rabbit mAb



Orders ■ 877-616-CELL (2355)
orders@cellsignal.com
Support ■ 877-678-TECH (8324)
info@cellsignal.com
Web ■ www.cellsignal.com

rev. 01/07/16

For Research Use Only. Not For Use In Diagnostic Procedures.

Applications	Species Cross-Reactivity*	Molecular Wt.	Isotype
W, IP Endogenous	H	80 kDa (NPM-ALK) 220 kDa (ALK)	Rabbit IgG**

Background: Anaplastic lymphoma kinase (ALK) is a tyrosine kinase receptor for pleiotrophin (PTN), a growth factor involved in embryonic brain development (1-3). In ALK-expressing cells, PTN induces phosphorylation of both ALK and the downstream effectors IRS-1, Shc, PLCγ, and PI3 kinase (1). ALK was originally discovered as a nucleophosmin (NPM)-ALK fusion protein produced by a translocation (4). Investigators have found that the NPM-ALK fusion protein is a constitutively active, oncogenic tyrosine kinase associated with anaplastic lymphoma (4). Research literature suggests that activation of PLCγ by NPM-ALK may be a crucial step for its mitogenic activity and involved in the pathogenesis of anaplastic lymphomas (5).


A distinct ALK oncogenic fusion protein involving ALK and echinoderm microtubule-associated protein like 4 (EML4) has been described in the research literature from a non-small cell lung cancer (NSCLC) cell line, with corresponding fusion transcripts present in some cases of lung adenocarcinoma. The short, amino-terminal region of the microtubule-associated protein EML4 is fused to the kinase domain of ALK (6-8).

Specificity/Sensitivity: ALK (C26G7) Rabbit mAb detects endogenous levels of total ALK protein. This antibody does not cross-react with other family members.

Source/Purification: Monoclonal antibody is produced by immunizing animals with a recombinant fusion protein surrounding amino acid 1475 of human ALK.

Background References:

- (1) Stoica, G.E. et al. (2001) *J Biol Chem* 276, 16772-9.
- (2) Iwahara, T. et al. (1997) *Oncogene* 14, 439-49.
- (3) Morris, S.W. et al. (1997) *Oncogene* 14, 2175-88.
- (4) Morris, S.W. et al. (1994) *Science* 263, 1281-4.
- (5) Bai, R.Y. et al. (1998) *Mol Cell Biol* 18, 6951-61.
- (6) Rikova, K. et al. (2007) *Cell* 131, 1190-203.
- (7) Takeuchi, K. et al. (2008) *Clin Cancer Res* 14, 6618-24.
- (8) Soda, M. et al. (2007) *Nature* 448, 561-6.



Western blot analysis of extracts from Karpas-299 and Sup-M2 cells using ALK (C26G7) Rabbit mAb. Cell Line Source: Dr Abraham Karpas at the University of Cambridge.

Entrez-Gene ID #238
UniProt ID #Q9UM73

Storage: Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

*Species cross-reactivity is determined by western blot.

**Anti-rabbit secondary antibodies must be used to detect this antibody.

Recommended Antibody Dilutions:

Western blotting 1:1000
Immunoprecipitation 1:200

For application specific protocols please see the web page for this product at www.cellsignal.com.

Please visit www.cellsignal.com for a complete listing of recommended companion products.

IMPORTANT: For western blots, incubate membrane with diluted antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

U.S. Patent No. 5,675,063

Tween® is a registered trademark of ICI Americas, Inc.