



**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Store at -20C  
#3347

## Synaptotagmin-1 Antibody

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W	H M R Mk	Endogenous	60	Rabbit	#P21579	6857

### Product Usage Information

#### Application

Western Blotting

#### Dilution

1:1000

### Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

### Specificity/Sensitivity

Synaptotagmin-1 Antibody detects endogenous levels of total synaptotagmin-1 protein.

### Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to human synaptotagmin-1 protein. Antibodies are purified by protein A and peptide affinity chromatography.

### Background

Synaptotagmin-1 (SYT1) is an integral membrane protein found in synaptic vesicles thought to play a role in vesicle trafficking and exocytosis (1). Individual SYT1 proteins are composed of an amino-terminal transmembrane region, a central linker region and a pair of carboxy-terminal C2 domains responsible for binding Ca<sup>2+</sup> (2). The C2 domains appear to be functionally distinct, with the C2A domain responsible for regulating synaptic vesicle fusion in a calcium-dependent manner during exocytosis while the C2B domain allows for interaction between adjacent SYT1 proteins (3). Because synaptotagmin 1 binds calcium and is found in synaptic vesicles, this integral membrane protein is thought to act as a calcium sensor in fast synaptic vesicle exocytosis. Evidence suggests possible roles in vesicle-mediated endocytosis and glucose-induced insulin secretion as well (4,5). SYT1 binds several different SNARE proteins during calcium-mediated vesicle endocytosis and an association between SYT1 and the SNARE protein SNAP-25 is thought to be a key element in vesicle-mediated exocytosis (6).

### Background References

1. Fukuda, M. and Mikoshiba, K. (2001) *Biochem Biophys Res Commun* 281, 1226-33.
2. Südhof, T.C. (2002) *J Biol Chem* 277, 7629-32.
3. Fernández-Chacón, R. et al. (2001) *Nature* 410, 41-9.
4. Lynch, K.L. et al. (2007) *Mol Biol Cell* 18, 4957-68.
5. Gauthier, B.R. and Wollheim, C.B. (2008) *Am J Physiol Endocrinol Metab* 295, E1279-86.
6. Bai, J. et al. (2004) *Neuron* 41, 929-42.

### Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

### Western Blot Buffer

**IMPORTANT:** For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

### Applications Key

**W:** Western Blotting

### Cross-Reactivity Key

**H:** Human **M:** Mouse **R:** Rat **Mk:** Monkey

### Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

### Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any

purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.