



**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Store at -20C  
#3455

## Phospho-DRP1 (Ser616) Antibody

For Research Use Only. Not for Use in Diagnostic Procedures.

<b>Applications:</b> W, W-S, IP, IF-IC, FC-FP	<b>Reactivity:</b> H	<b>Sensitivity:</b> Endogenous	<b>MW (kDa):</b> 78-82	<b>Source/Isotype:</b> Rabbit	<b>UniProt ID:</b> #O00429	<b>Entrez-Gene Id:</b> 10059
--	-------------------------	-----------------------------------	---------------------------	----------------------------------	-------------------------------	---------------------------------

### Product Usage Information

#### Application

Western Blotting  
Simple Western™  
Immunoprecipitation  
Immunofluorescence (Immunocytochemistry)  
Flow Cytometry (Fixed/Permeabilized)

#### Dilution

1:1000  
1:10 - 1:50  
1:50  
1:200 - 1:400  
1:50 - 1:200

### Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

### Specificity/Sensitivity

Phospho-DRP1 (Ser616) Antibody detects endogenous levels of DRP1 only when phosphorylated at Ser616.

### Species predicted to react based on 100% sequence homology

Mouse, Rat, Monkey

### Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Ser616 of human DRP1. Antibodies are purified by protein A and peptide affinity chromatography.

### Background

Dynamin-related protein 1 (DRP1) is a member of the dynamin superfamily of GTPases. Members of this family have diverse cellular functions including vesicle scission, organelle fission, viral resistance, and intracellular trafficking (reviewed in 1). DRP1 affects mitochondrial morphology and is important in mitochondrial and peroxisomal fission in mammalian cells (2-5). The yeast ortholog of DRP1 clusters into a spiral-shaped structure on the mitochondrial membrane at the site of fission (reviewed in 6), and this structure is likely conserved in mammalian cells (3). The division of the mitochondria, which is required for apoptosis, as well as normal cell growth and development is controlled, in part, by the phosphorylation of DRP1 at Ser616 by Cdk1/cyclin B and at Ser637 by protein kinase A (PKA) (reviewed in 6). When phosphorylated at Ser616, DRP1 stimulates mitochondrial fission during mitosis. Conversely, fission is inhibited when DRP1 is phosphorylated at Ser637 (reviewed in 6). Dephosphorylation at Ser637 by calcineurin reverses this inhibition (7). In addition to phosphorylation, sumoylation of DRP1 is also an enhancer of mitochondrial fission (8). Balancing fission and fusion events is essential for proper mitochondrial function. Research studies have demonstrated mitochondrial defects in a variety of neurodegenerative diseases including Alzheimer's disease, Parkinson's disease, and Huntington's disease (reviewed in 6).

### Background References

1. Praefcke, G.J. and McMahon, H.T. (2004) *Nat Rev Mol Cell Biol* 5, 133-47.
2. Taguchi, N. et al. (2007) *J Biol Chem* 282, 11521-9.
3. Smirnova, E. et al. (2001) *Mol Biol Cell* 12, 2245-56.
4. Smirnova, E. et al. (1998) *J Cell Biol* 143, 351-8.
5. Koch, A. et al. (2003) *J Biol Chem* 278, 8597-605.
6. Knott, A.B. et al. (2008) *Nat Rev Neurosci* 9, 505-18.
7. Cereghetti, G.M. et al. (2008) *Proc Natl Acad Sci USA* 105, 15803-8.
8. Zunino, R. et al. (2007) *J Cell Sci* 120, 1178-88.

### Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

### Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

**Applications Key**

**W:** Western Blotting **W-S:** Simple Western™ **IP:** Immunoprecipitation **IF-IC:** Immunofluorescence (Immunocytochemistry) **FC-FP:** Flow Cytometry (Fixed/Permeabilized)

**Cross-Reactivity Key**

**H:** Human

**Trademarks and Patents**

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

**Limited Uses**

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.