



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Store at -20C
#3484

TRPV3 Antibody

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W, IP	H M	Endogenous	98	Rabbit	#Q8NET8	162514
Product Usage Information	Application			Dilution		
	Western Blotting			1:1000		
	Immunoprecipitation			1:50		
Storage	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.					
Specificity/Sensitivity	TRPV3 Antibody detects endogenous levels of total TRPV3 protein.					
Source / Purification	Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Asn101 of human TRPV3. Antibodies are purified by peptide affinity chromatography.					
Background	The transient receptor potential cation channel subfamily V member 3 (TRPV3 or VRL3) is a widely expressed ligand-gated ion channel protein that acts in temperature detection between 22° C and 40° C (1). Related transient receptor proteins (TRP) respond to different temperature ranges; TRPA1 responds to cold temperatures and TRPM1 to cool, while warmer temperatures elicit response from TRPV3 and TRPV4. Noxious heat or painful temperatures correlate with TRPM8 and TRPV1 activity (reviewed in 2). TRPV3 is found in association with TRPV1 (VR1) and is thought to modulate activity of this noxious heat-responsive protein (3). Increased TRPV1 and TRPV3 protein expression is correlated with nervous injury, while reduced expression of both may be seen in cases of diabetic skin neuropathy (2). Recent evidence implicates TRPV3 in pruritic dermatitis, or itching associated with eczema and related skin disorders (4).					
Background References	<ol style="list-style-type: none"> Xu, H. et al. (2002) <i>Nature</i> 418, 181-6. Facer, P. et al. (2007) <i>BMC Neurol</i> 7, 11. Smith, G.D. et al. (2002) <i>Nature</i> 418, 186-90. Steinhoff, M. and Bíró, T. (2009) <i>J Invest Dermatol</i> 129, 531-5. 					

Species Reactivity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).
Western Blot Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.
Applications Key	W: Western Blotting IP: Immunoprecipitation
Cross-Reactivity Key	H: Human M: Mouse
Trademarks and Patents	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.
Limited Uses	Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect. Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for

diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.