

Store at -20C  #3523	RalB Antibody	
		<b>Orders:</b> 877-616-CELL (2355) orders@cellsignal.com
		<b>Support:</b> 877-678-TECH (8324)
		<b>Web:</b> info@cellsignal.com cellsignal.com
3 Trask Lane   Danvers   Massachusetts   01923   USA		

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W	H Mk	Endogenous	26	Rabbit	#P11234	5899
<b>Product Usage Information</b>	<b>Application</b>		<b>Dilution</b>			
	Western Blotting		1:1000			
<b>Storage</b>	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.					
<b>Specificity/Sensitivity</b>	RalB Antibody detects endogenous levels of total RalB protein. The antibody might also react with overexpressed RalA protein.					
<b>Source / Purification</b>	Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Glu142 of human RalB protein. Antibodies are purified by peptide affinity chromatography.					
<b>Background</b>	RalA and RalB are members of the Ras family of small GTPases and are highly homologous in protein sequence. The functions of RalA and RalB are distinct yet overlapping. By binding to various effector proteins, RalA and RalB serve as important GTP sensors for exocytosis and membrane trafficking (1-3). RalA is required for Ras-related tumorigenesis (4) and RalB is important for tumor survival (5). In addition to tumor formation, Ral proteins also play a role in cancer cell migration and metastatic tumor invasion (6,7).					
<b>Background References</b>	<ol style="list-style-type: none"> <li>1. Li, G. et al. (2007) <i>J Neurosci</i> 27, 190-202.</li> <li>2. van Dam, E.M. and Robinson, P.J. (2006) <i>Int J Biochem Cell Biol</i> 38, 1841-7.</li> <li>3. Camonis, J.H. and White, M.A. (2005) <i>Trends Cell Biol</i> 15, 327-32.</li> <li>4. Lim, K.H. et al. (2005) <i>Cancer Cell</i> 7, 533-45.</li> <li>5. Chien, Y. et al. (2006) <i>Cell</i> 127, 157-70.</li> <li>6. Oxford, G. et al. (2005) <i>Cancer Res</i> 65, 7111-20.</li> <li>7. Lim, K.H. et al. (2006) <i>Curr Biol</i> 16, 2385-94.</li> </ol>					

**Species Reactivity** Species reactivity is determined by testing in at least one approved application (e.g., western blot).

**Western Blot Buffer** IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween@ 20 at 4°C with gentle shaking, overnight.

**Applications Key** **W:** Western Blotting

**Cross-Reactivity Key** **H:** Human **Mk:** Monkey

**Trademarks and Patents** Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.  
All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

**Limited Uses** Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not

sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.

**Orders: 877-616-CELL (2355) • [orders@cellsignal.com](mailto:orders@cellsignal.com) • Support: 877-678-TECH (8324) • [info@cellsignal.com](mailto:info@cellsignal.com) • Web: [cellsignal.com](http://cellsignal.com)**  
For Research Use Only. Not for Use in Diagnostic Procedures.