Nod1 Antibody



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W	Reactivity: H M R Mk	Sensitivity: Endogenous	MW (kDa): 95	Source/Isotype: Rabbit	UniProt ID: #Q9Y239	Entrez-Gene Id: 10392
Product Usage Information		Application Western Blotting			Dilution 1:1000	
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA and 50% glycerol. Store at – 20°C. Do not aliquot the antibody.				
Specificity/Sensitivity		Nod1 Antibody detects endogenous levels of Nod1 protein.				
Source / Purification		Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues near the carboxy terminus of Nod1 protein. Antibodies were purified by protein A and peptide affinity chromatography.				
Background		Nod1/CARD4 is a cytosolic protein structually related to Apaf-1 and plant drug-resistance proteins that has been implicated in apoptosis and inflammatory responses to certain pathogenic bacteria (1-3). It contains an amino-terminal caspase recruitment domain (CARD) that is linked to a central nucleotide-binding domain (NBD; also known as a NOD domain) and is followed by carboxy-terminal leucine-rich repeats (LRR) (1). Like Apaf-1, Nod1 induces apoptosis by a CARD/NBD-dependent activation of caspase-9 (1). The primary function of Nod1 is thought to be as a sensor for certain pathogenic microbes and triggering inflammatory responses including the activation of NF-kB and JNK pathways (4-6). The LRR of Nod1 appears to be involved in recognition of microbial components and the CARD domain induces NF-kB activation in cooperation with the CARD containing kinase, RICK/RIP2/CARDIAK (1,5,6). Mutations in Nod1 have been linked increased susceptibility to asthma (7) and inflammatory bowel disease (8).				
Background References		1. Inohara, N. et al. (1999) <i>J. Biol. Chem.</i> 274, 14560-14567. 2. Inohara, N. and Nuñez, G. (2003) <i>Nat. Rev. Immunol.</i> 3, 371-382. 3. Fritz, J.H. et al. (2006) <i>Nat .Immunol.</i> 7, 1250-1257. 4. Girardin, S.E. et al. (2001) <i>EMBO Rep.</i> 2, 736-742. 5. Inohara, N. et al. (2001) <i>J. Biol. Chem.</i> 276, 2551-2554. 6. Inohara, N. et al. (2000) <i>J. Biol. Chem.</i> 275, 27823-27831. 7. Hysi, P. et al. (2005) <i>Hum. Mol. Genet.</i> 14, 935-941. 8. McGovern, D.P. et al. (2005) <i>Hum. Mol. Genet.</i> 14, 1245-1250.				

Species Reactivity Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X

TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key W: Western Blotting

Cross-Reactivity Key H: Human M: Mouse R: Rat Mk: Monkey

Trademarks and Patents Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for

more information.

Limited Uses Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST,

the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless

separately accepted in writing by a legally authorized representative of CST, are rejected and are of no

force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.