



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

CDX2 Antibody

Store at -20C
#3977

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, IP	Reactivity: H	Sensitivity: Endogenous	MW (kDa): 38	Source/Isotype: Rabbit	UniProt ID: #Q99626	Entrez-Gene Id: 1045
-------------------------------	-------------------------	-----------------------------------	------------------------	----------------------------------	-------------------------------	--------------------------------

Product Usage Information

Application

Western Blotting
Immunoprecipitation

Dilution

1:1000
1:100

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

CDX2 Antibody detects endogenous levels of total CDX2 protein.

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to amino acids at the N-terminus of human CDX2. Antibodies are purified by Protein A and peptide affinity chromatography.

Background

CDX2, a homeobox domain-containing transcription factor, is a master regulator of the trophectoderm, the layer that gives rise to extra-embryonic tissues in mammalian development (1). CDX2 is also involved in intestinal development (2), and gain of expression or loss of expression has been associated with various human malignancies, such as Barrett's esophagus (3) and colorectal cancer (4,5). Mouse embryonic stem cells deficient in CDX2 display limited hematopoietic progenitor development and altered *Hox* gene expression (6), pointing to a role for CDX2 in *Hox* gene regulation. CDX2 is also implicated in the aberrant expression of *Hox* genes in human AML cell lines (7).

Background References

1. Ralston, A. and Rossant, J. (2008) *Dev Biol* 313, 614-29.
2. Traber, P.G. and Silberg, D.G. (1996) *Annu Rev Physiol* 58, 275-97.
3. Shi, X.Y. et al. (2008) *Am J Clin Pathol* 129, 571-7.
4. Bakaris, S. et al. (2008) *Histol Histopathol* 23, 1043-7.
5. Guo, R.J. et al. (2004) *Cancer Biol Ther* 3, 593-601.
6. Wang, Y. et al. (2008) *Proc Natl Acad Sci U S A* 105, 7756-61.
7. Fröhling, S. et al. (2007) *Cell Cycle* 6, 2241-5.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween@ 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IP:** Immunoprecipitation

Cross-Reactivity Key

H: Human

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for

Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.