

**Limited Uses** 

## LIN28A (A177) Antibody



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

## For Research Use Only. Not for Use in Diagnostic Procedures.

<b>Applications:</b> W, IF-IC, FC-FP	Reactivity: H M	<b>Sensitivity:</b> Endogenous	<b>MW (kDa):</b> 26	Source/Isotype: Rabbit	UniProt ID: #Q9H9Z2	Entrez-Gene Id: 79727
Product Usage Information		Application Western Blotting Immunofluorescence Flow Cytometry (Fixed	•	istry)		<b>Dilution</b> 1:1000 1:800 1:100
Storage	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA and 5 20°C. Do not aliquot the antibody.				/ml BSA and 50% g	lycerol. Store at –
Specificity/Sensitivity		LIN28A (A177) Antibody detects endogenous levels of LIN28A protein.				
Species predicted to react based on 100% sequence homology		Monkey				
Source / Purification		Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to amino acid sequence surrounding Ala177 of human LIN28A. Antibodies are purified by protein A and peptide affinity chromatography.				
Background		LIN28A and LIN28B are conserved, developmentally regulated RNA binding proteins that inhibit the processing and maturation of the let-7 family of miRNAs (1,2). The let-7 miRNAs have been implicated in repression of oncogenes such as Ras, Myc, and HMGA2 (3). It has recently been shown that upregulation of LIN28A and LIN28B in primary human tumors and human cancer cell lines is correlated with downregulation of let-7 miRNAs (4). LIN28 genes are reported to be involved in primordial germ cell development and germ cell malignancy (5). In addition, allelic variation in LIN28B is associated with regulating the timing of puberty in humans (6). Overexpression of LIN28A, in conjunction with Oct-4, Sox2, and Nanog, can reprogram human fibroblasts to pluripotent, ES-like cells (7).				
Background References		<ol> <li>Balzer, E. and Moss, E.G. (2007) RNA Biol 4, 16-25.</li> <li>Piskounova, E. et al. (2008) J Biol Chem 283, 21310-4.</li> <li>Cho, W.C. (2007) Mol Cancer 6, 60.</li> <li>Viswanathan, S.R. et al. (2009) Nat Genet 41, 843-8.</li> <li>West, J.A. et al. (2009) Nature 460, 909-13.</li> <li>Ong, K.K. et al. (2009) Nat Genet 41, 729-33.</li> <li>Yu, J. et al. (2007) Science 318, 1917-20.</li> </ol>				
Species Reactivit	ty	Species reactivity is de	etermined by testin	g in at least one approve	ed application (e.g.,	western blot).
Western Blot Buffer		IMPORTANT: For western blots, incubate membrane with diluted primary antibody in TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.				n 5% w/v BSA, 1X
Applications Key		<b>W:</b> Western Blotting <b>IF-IC:</b> Immunofluorescence (Immunocytochemistry) <b>FC-FP:</b> Flow (Fixed/Permeabilized)				w Cytometry
Cross-Reactivity Key		H: Human M: Mouse				
Trademarks and Patents		Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.				

more information.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's

terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.